

In-vivo anatomical reconstruction of the optic radiations in the human brain

Demetrio Milardi^{1,2}, Fabio Trimarchi¹, Alessandro Arrigo^{1,3}, Alessandro Calamuneri¹, Angelo Quartarone¹, Giuseppe Santoro¹, Giuseppina Rizzo¹, Grazia Giulia Pantè¹, Alba Migliorato¹, Salvatore Bertino¹

¹Department of Biomedical, Dental Sciences and Morphological and Functional Images, University of Messina, 98125 Messina, Italy

²IRCCS Centro Neurolesi "Bonino Pulejo", Messina, Italy

³Department of Ophthalmology, University Vita e Salute San Raffaele, Milano, Italy

The optic radiations are major white matter pathways funneling visual information from the lateral geniculate nuclei to the visual cortex in the occipital lobe. Given their relevance in visual processing and in several brain disorders, the optic radiations have been extensively investigated by using magnetic resonance imaging and diffusion tensor imaging tractography [1]. Herein, we use a powerful diffusion signal modeling, namely Constrained Spherical Deconvolution, in order to provide an exhaustive connectivity profile of the connections between the lateral geniculate nucleus and the visual cortex in the healthy brain, as well as pulvinar connectivity with visual-related structures. In addition, taking into account that visual deficits may precede motor symptoms' onset in Parkinson's Disease [2], we assessed whether the intracranial visual system can be involved at the early stage of the disease. Our connectivity analysis revealed that the optic radiations are mainly distributed in V1 and V2. Furthermore, we found significant alterations of optic radiations connectivity distribution in Parkinson's Disease patients, with decreased lateral geniculate nucleus-V2 density as well as significant increase of optic radiations' mean diffusivity. Voxel Based Morphometry analysis also showed significant reduction of visual cortical volumes and of the optic radiation in the patients group. In conclusion, our findings provide a reliable connectivity profile of the optic radiations, suggesting extrastriate-lateral geniculate nucleus connections in human brain. Finally, we showed that visual system alterations can be detected at early stages of Parkinson's Disease.

References

- [1] Kuhnt et al (2013). Optic radiation fiber tractography in glioma patients based on high angular resolution diffusion imaging with compressed sensing compared with diffusion tensor imaging - initial experience. *PLoS One* 8: e70973.
- [2] Bioussé et al (2004). Ophthalmologic features of Parkinson's disease. *Neurology* 62:177-180.

Keywords

Optic radiations, tractography, MRI, Parkinson's Disease, brain connectivity