

## Hepatic matrix metalloproteinase-10 exerts a hepatoprotective role after acute liver injury

Simone Carotti<sup>1</sup> - Oihane Garcia-Irigoyen<sup>2</sup> - Maria U. Latasa<sup>2</sup> - Iker Uriarte<sup>2</sup> - Maite G. Fernandez-Barrena<sup>2</sup> - Maria Elizalde<sup>2</sup> - Raquel Urtasun<sup>2</sup> - Umberto Vespasiani-Gentilucci<sup>3</sup> - Jesus M. Banales<sup>4</sup> - William C. Parks<sup>5</sup> - Jose A. Rodriguez<sup>6</sup> - Josune Orbe<sup>6</sup> - Jesus Prieto<sup>2</sup> - Jose A. Paramo<sup>6</sup> - Carmen Berasain<sup>2</sup> - Matias A. Avila<sup>2</sup> - Sergio Morini<sup>1</sup>

<sup>1</sup>Unit of Microscopic and Ultrastructural Anatomy, Department of Medicine, Università Campus Bio-Medico di Roma, Roma, Italia - <sup>2</sup>Hepatology Program, Centro de Investigación Médica Aplicada (CIMA), University of Navarra, Pamplona, Spagna - <sup>3</sup>Unit of Internal Medicine and Hepatology, Department of Medicine, Università Campus Bio-Medico di Roma, Roma, Italia - <sup>4</sup>Department of Liver and Gastrointestinal Diseases, Biodonostia Research Institute-Donostia University Hospital, University of Basque Country, San Sebastian, Spagna - <sup>5</sup>Center for Lung Biology, University of Washington, Seattle, Stati Uniti d'America - <sup>6</sup>Centro de Investigación Médica Aplicada (CIMA), Division of Cardiovascular Sciences, Universidad de Navarra, Pamplona, Spain.

After injuries that lead to a loss of liver tissue a regenerative and reparative response is performed in order to restore an adequate hepatic mass. The remodeling of the extracellular matrix, accompanies the liver regeneration and when the reparative reaction goes awry in the setting of chronic liver injury, could be involved in the carcinogenic process (1,2). Following the damage, a provisional matrix is deposited, intended to be successively replaced, which has the function of stabilizing the lesional area and constitutes a support for guiding regenerating cells. Matrix metalloproteinases are increasingly recognized as important modulators of the matrix remodeling process. Matrix metalloproteinase-10 (MMP-10) has been implicated in the reparative process in other organs and has effects on the plasminogen system, which plays a fundamental role in liver repair (3). The hepatic expression of MMP10 in animal models of acute liver injury was tested in order to investigate the role of MMP-10 in liver repair and regeneration. The liver regeneration after two thirds partial hepatectomy (PH) and bile duct ligation (BDL) models were examined. Hepatic MMP-10 expression, analyzed by immunohistochemistry, western blot and qPCR showed a rise early after injury. In the MMP10-deficient mice a diminished and delayed resolution of necrotic lesions, enhanced fibrogenesis and a fibrinogen/fibrin and fibronectin compromised turnover were observed. These findings showed that the MMP10 expression plays a role in the hepatic wound healing response probably through its profibrinolytic activity.

### References

- [1] Uriarte et al. (2013) Identification of fibroblast growth factor 15 as a novel mediator of liver regeneration and its application in the prevention of post-resection liver failure in mice. *Gut*;62(6):899-910.
- [2] Uriarte et al. (2015) Ileal FGF15 contributes to fibrosis-associated hepatocellular carcinoma development. *Int J Cancer*; 136(10):2469-75.
- [3] Orbe et al. (2011) Matrix metalloproteinase-10 effectively reduces infarct size in experimental stroke by enhancing fibrinolysis via a thrombin-activatable fibrinolysis inhibitor-mediated mechanism. *Circulation*; 124(25):2909-19.

### Key words

Metalloproteinase-10; liver repair and regeneration; hepatocarcinogenesis.