

S100B protein regulates myoblast and macrophage functions in skeletal muscle regeneration

Francesca Riuzzi - Sara Beccafico - Roberta Sagheddu - Sara Chiappalupi - Ileana Giambanco - Guglielmo Sorci - Rosario Donato

Università degli Studi di Perugia, Dipartimento di Medicina Sperimentale, Perugia, Italia

Regeneration of acutely injured skeletal muscles relies on a tightly controlled chain of cellular and molecular events, but a complete picture of factors concurring to the regeneration process is still missing. Extracellular S100B protein inhibits myoblast differentiation and stimulates myoblast proliferation by activating its canonical receptor, RAGE (receptor for advanced glycation endproducts), or bFGF/FGFR1 depending on myoblast density (1-4). S100B is released by damaged muscle tissue early after injury in advance of bFGF release, with declining release thereafter (4). We show that S100B is required for correct timing of skeletal muscle regeneration after acute injury. S100B expands the myoblast population, attracts macrophages to damage sites, promotes macrophage polarization into M2 (pro-regenerative) phenotype and reduces fibroblast proliferation. Also, S100B is transiently induced in and released by infiltrating macrophages under the action of proinflammatory and antiinflammatory cytokines, and effects of macrophage-derived S100B sum up with those of myofiber-released S100B. S100B's effects are mediated by RAGE during the first 3 days after injury, however during the myoblast proliferation phase/macrophage M2 phase (i.e. at days 4-6 post-injury) S100B also activates bFGF-FGFR1 to stimulate myoblast proliferation and macrophage M1/M2 transition. Thus, S100B is a major molecular determinant of timed muscle regeneration after acute injury by virtue of its regulatory effects on myoblasts and macrophages.

This work was supported by grants from MIUR PRIN-2010R8JK2X_004, AFM-Téléthon 16260 and Fondazione CRP 2012.0241.021.

References

- [1] Sorci et al. (2003) S100B inhibits myogenic differentiation and myotube formation in a RAGE-independent manner. *Mol Cell Biol* 23, 4870-4881.
- [2] Riuzzi et al. (2006) S100B stimulates myoblast proliferation and inhibits myoblast differentiation by independently stimulating ERK1/2 and inhibiting p38 MAPK. *J Cell Physiol* 207, 461-470.
- [3] Riuzzi et al. (2011) S100B protein regulates myoblast proliferation and differentiation by activating FGFR1 in a bFGF-dependent manner. *J Cell Sci* 124, 2389-2400, doi: 10.1242/jcs.084491.
- [4] Riuzzi et al. (2012) S100B Engages RAGE or bFGF/FGFR1 in Myoblasts Depending on Its Own Concentration and Myoblast Density. Implications for Muscle Regeneration. *PLoS ONE* 7(1): e28700. doi: 10.1371/journal.pone.0028700.

Keywords

Muscle regeneration; S100B; RAGE; bFGF/FGFR1; myoblasts; macrophages.