

Effect of hypoxia-inducing ions on chondrogenic differentiation in adipose derived mesenchymal stem cells within alginate matrix

Stefano Focaroli¹ - Viviana Salvatore¹ - Nicola Zanna² - Alessandra Ruggeri¹ - Antonio Mazzotti³ - Claudia Tomasini²

¹Universita' di Bologna, Dipartimento di Scienze Biomediche e Neuromotorie, Bologna, Italia - ²Universita' di Bologna, Dipartimento di Chimica "Giacomo Ciamician", Bologna, Italia - ³Universita' di Bologna, Dipartimento di Scienze Biomediche e Neuromotorie, Istituto Ortopedico Rizzoli, Bologna, Italia

Cartilage is a highly organized tissue with complex biomechanical properties, but since it has a poor intrinsic capacity of self-healing, injuries at this site usually lead to several problems, often ending in disabling symptoms. Although, different approaches have been proposed, even now cartilage repair represents a great challenge for orthopaedic surgeons (1, 2). One of the promising approach is given from tissue engineering, employing the combination of biomaterials and cell therapy to develop new therapeutic strategies. In this paper, we describe the behaviour of human adipose derived mesenchymal stem cells encapsulated into Ca/Co alginate beads as potential chondrogenic inducing biomaterial tacking advance on the synergy between alginate matrix and Co+2 ions without employing other expensive growth factors such as TGFbs or BMPs. The expression of chondrogenic markers such as sox9, collagen type II, and versican was investigated by Real Time PCR and Western blotting analysis. The expression of hif1mRNA was investigated to check the capability of Co+2 ions to induce a chemical hypoxia. Real Time PCR and WB data reveal a different cells behaviour on chondrogenic marker expression. In conclusion, the synergic effect of alginate and Co+2 ions can represent a valid strategy for chondrogenic differentiation of stem cells.

References

- [1] Ling et al. (2013) Regeneration of Articular Cartilage by Adipose Tissue Derived Mesenchymal Stem Cells: Perspectives From Stem Cell Biology and Molecular Medicine. *J. Cell. Physiol.* 228: 938.
- [2] Wei et al. (2012) Chondrogenic differentiation of induced pluripotent stem cells from osteoarthritic chondrocytes in alginate matrix. *Eur. Cell. Mat.* 23: 1.

Keywords

Stem cells; chondrogenic differentiation; alginate.