

From interaction to function: Phospholipase C beta 1 protects cells from stress-induced apoptosis

Manuela Piazzì¹, Alberto Bavelloni^{2,3}, William Blalock^{2,4}, Irene Faenza¹

¹ Cell Signaling Laboratory, Department of Biomedical Science, University of Bologna, IT

² SC Laboratory of Musculoskeletal Cell Biology, Rizzoli Orthopedic Institute, Bologna, Italy

³ Laboratory RAMSES, Rizzoli Orthopedic Institute, Bologna, Italy

⁴ Institute of Molecular Genetics, National Research Council of Italy (IGM-CNR), Bologna, Italy

The phosphoinositide-dependent signal transduction pathway has been implicated in the control of a variety of biological processes, such as the regulation of cellular metabolism and homeostasis, cell proliferation and differentiation. One of the key player in the regulation of inositol lipid signaling is phospholipase C beta 1 (PI-PLC β 1), which hydrolyses PtdIns(4,5)P₂, giving rise to the second messengers IP₃ and DAG.

The complete mapping of the PI-PLC β 1 interactome was undertaken, to understand its diverse functions within the nuclear compartment and to determine its contribution to physiological and pathological processes. Affinity purification-mass spectrometry (AP-MS) allowed for the identification of 160 proteins present in association with PI-PLC β 1 in the nucleus of erythroleukemia cells. Co-immunoprecipitation analysis of selected proteins confirmed the data obtained from mass spectrometry. Of particular interest was the identification of proteins involved in nuclear trafficking, as well as factors involved in hematological malignancies and several anti-apoptotic proteins (Piazzì et al., 2013).

PI-PLC β 1 has been associated with the regulation of several cellular functions, some of which are not yet fully understood. In particular, it has been reported that PI-PLC β 1 protects murine fibroblasts from oxidative stress-induced cell death, through signaling events which remain unclear. Reactive oxygen species (ROS) have been shown to regulate major epigenetic processes causing the silencing of tumor suppressors and enhancing the proliferation of leukemic cells under oxidative stress. Investigation of the role for ROS and their signaling mediators in the pathogenesis of leukemia might, therefore, outline innovative approaches for the improvement of pharmacological therapies to treat leukemia.

We demonstrate that in acute lymphoid leukemia cells (pro-B cells), treated with 250 μ M of hydrogen peroxide (H₂O₂), PI-PLC β 1b conferred resistance to cell death, promoting cell cycle progression and cell proliferation. Interestingly, we found that, upon H₂O₂ exposure, the expression of PI-PLC β 1b affects the activity of several protein kinases, in particular it completely abolished the phosphorylation of Erk1/2 MAP kinases, down-regulated PTEN and up-regulated the phosphorylation of Akt; thereby sustaining cellular proliferation.

Reference

- [1] Piazzì M et al. (2013). Phosphoinositide-specific phospholipase C β 1b (PI-PLC β 1b) interactome: affinity purification-mass spectrometry analysis of PI-PLC β 1b association with nuclear proteins. *Mol Cell Prot* 12:2220-35.