

## Hypoxia induces Galectin-3 and Bcl-2 over-expression in human umbilical vein endothelial cells (HUVECs)

Anna Di Vito<sup>1</sup>, Chiara Mignogna<sup>2</sup>, Anna Bilotta<sup>1</sup>, Caterina Camastra<sup>2</sup>, Francesco Trapasso<sup>1</sup>, Antonio Brunetti<sup>3</sup>, Ilenia Aversa<sup>1</sup>, Giuseppe Donato<sup>2</sup>, Tullio Barni<sup>1</sup>

<sup>1</sup> Clinical and Experimental Medicine Department, University of Catanzaro "Magna Græcia", Catanzaro, Italy

- <sup>2</sup> Health Science Department, Pathology Unit, University of Catanzaro "Magna Græcia", Catanzaro, Italy -

<sup>3</sup> Health Science Department, University of Catanzaro "Magna Græcia", Catanzaro, Italy

Angiogenesis, the growth of new blood vessels from pre-existing endothelium, is a critical phenomenon occurring during development and tissue regeneration. In pathological conditions such as inflammation and malignancies, hypoxia represent one of the most important driver of angiogenesis, mainly via the release of nitric oxide [1]. Here we investigate the behavior of human umbilical vein endothelial cells (HUVECs) treated with 100 mM CoCl<sub>2</sub> for 24 hours, a condition mimicking hypoxia by the stabilization of HIF-1 $\alpha$  and HIF-2 $\alpha$  [2,3]. MTT and wound healing assays were performed to evaluate cell migration and proliferation, respectively, while Bcl-2 and Galectin-3 expression levels were analyzed by western blotting. We showed that hypoxic condition resulted in reduced proliferation and migration with increased expression of Galectin-3 and Bcl-2. These preliminary results provide new insights in the characterization of Galectin-3/Bcl-2 interplay in endothelial cell survival under hypoxic condition, and will contribute to a better understanding of hypoxia influences on tumor angiogenesis.

### References

- [1] Wang Z et al. (2015) Broad targeting of angiogenesis for cancer prevention and therapy. *Semin Cancer Biol* pii: S1044-579X(15)00002-4; doi: 10.1016/j.semcancer.2015.01.001.
- [2] Ikemori RY et al. (2014) Galectin-3 Up-Regulation in Hypoxic and Nutrient Deprived Microenvironments Promotes Cell Survival. *PLoS One* 9(11): e111592; doi: 10.1371/journal.pone.0111592.
- [3] Doi N et al. (2015) The acquired radioresistance in HeLa cells under conditions mimicking hypoxia was attenuated by a decreased expression of HIF subunit genes induced by RNA interference. *Exp Cell Res* 333(2): 249-60; doi: 10.1016/j.yexcr.2015.03.009.

### Keywords

HUVEC; hypoxia; Galectin-3; Bcl-2.