

Potential protective effects of melatonin against UV-A irradiation on fibroblast cell line

Giovanni Damiani, Michele Peroni, Antonia Infranco and Francesca Bonomini

Human Anatomy Division, Department of Biomedical Science and Biotechnology, University of Brescia, Brescia, Italy

The sun's radiation that reaches Earth contains ultraviolet (UV) wavelights made up of a combination of UV-A (95%) and UV-B (5%) radiations. Chronic sun exposure is responsible for long term clinical skin changes such as photoaging, photodamage and photocancers. Moreover, inflammation is mostly due to UV-A which stimulates the production of reactive oxygen species (ROS) inducing also photoaging (Mouchet et al., 2010; Marionnet et al., 2010). In order to protect themselves against oxidative stress, skin cells developed several defense systems, including ROS and metal ions scavengers and a battery of detoxifying and repair enzymes (Bickers and Athar, 2006). In addition, UV-A can also directly influence the structure of nucleic acids, breaking the chain or changing the nucleotide sequence. Altogether these perturbations of cells homeostasis advantages a significant up-regulation of oxidative and inflammatory responsive genes. In this study, we focused our attention on prevention of photodamage, choosing melatonin as antioxidant agent. Melatonin is a neuroendocrine mediator with pleiotropic bioactivities such as hormonal, neurotransmitter, immunomodulator and biological modifier actions. Its antioxidant activity is the result of two different but synergic actions: a direct, due to its ability to act as a free radical scavenger and an indirect due to the up-regulation of antioxidant enzymes (Fischer et al., 2008). The aim of the present study was to analyze the impact of pre-treatment of murine fibroblasts cells (NIH-3T3) with melatonin (10⁻³ M- kindly provided by Chronolife S.r.l., Roma, Italy) later irradiated by UV-A irradiation (15 J/cm²) evaluating the changes of fibroblast microenvironment conditions. We observed that UV-A irradiation caused matrix restructuration and alteration, oxidative stress and inflammation; while melatonin pre-treatment suppresses UV-A induced photodamage. Collectively, these results suggest that melatonin provides relevant protective effects against UV-A irradiation. A new chapter of melatonin in dermato-endocrine research could be open.

References

- Mouchet et al. (2010) In vivo identification of Solar radiation-responsive gene network: role of the p38 stress-dependent kinase. PLoS ONE 5: e10776.
- Marionnet et al. (2010) Different Oxidative Stress Response in Keratinocytes and Fibroblasts of Reconstructed Skin Exposed to Non Extreme Daily-Ultraviolet Radiation. PLoS ONE 5: e12059.
- Bickers and Athar (2006) Oxidative stress in the pathogenesis of skin disease. J Invest Dermatol 126: 2565-2575.
- Fischer et al. (2008). Melatonin as a major skin protectant: from free radical scavenging to DNA damage repair. Exp Dermatol 17: 713-730.

Keywords: Fibroblast, melatonin, oxidative stress, UV-A irradiation.