

Analysis of tight junctions in placentas affected by chorioamnionitis: in vivo and in vitro analysis

Giovanni Tossetta¹, Francesca Paolinelli¹, Teresa Lorenzi¹, Pasquapina Ciarmela¹, Chiara Avellini¹, Eleonora Salvolini², Paolo Toti³, Felice Petraglia⁴, Mario Castellucci¹ and Daniela Marzioni¹

¹ Department of Experimental and Clinical Medicine Università Politecnica delle Marche, Ancona, Italy

² Department of Clinical and Molecular Sciences, Università Politecnica delle Marche, Ancona, Italy

³ Department of Human Pathology and Oncology University of Siena, Siena, Italy

⁴ Department of Pediatrics, Obstetrics and Reproductive Medicine University of Siena, Siena, Italy

The human placenta and fetal membranes provide a barrier regulating the transfer of materials between the mother and the developing fetus throughout gestation. Chorioamnionitis is an important risk factor for preterm delivery that is associated with high perinatal morbidity and mortality. Chorioamnionitis is the term applied to infections of the placenta and membranes resulting in high concentrations of IL-1beta, IL-6, IL-8 and TGF-beta in the amniotic fluid (D'Alquen et al., 2005). With progression of inflammation, immune cells penetrate blood vessels and infiltrate the umbilical cord, resulting in funisitis (Romero and Mazor, 1988). In normal conditions the two important physical entities in endothelial/epithelial paracellular clefts are adherens junctions and tight junctions.

Tight junction governs the paracellular movement of water, solutes and immune cells, through the intercellular space creating a boundary between the apical and basolateral sides of cellular barriers (Gruenheid and Finlay, 2003). We have evaluated the localization of tight junctions studying the Zonula Occludens-1 (ZO-1) and Occludin expressions as well as the localization of adherent junctions, testing the expression of VE-cadherin and beta-catenin in placentas from normal gestations, from preterm idiopathic deliveries and from chorioamnionitis by immunohistochemistry. In addition, we have evaluated the mRNAs by real time PCR, the protein levels of these molecules by Western blot analysis in placental tissues, and to better clarify the action of some cytokines on occludin we performed in vitro analysis of HUVEC cultures.

Our more striking result is the decrease of occludin expression in placentas from chorioamnionitis and an evident action of the cytokines on this molecule.

References

- D'Alquen et al. (2005) Activation of umbilical cord endothelial cells and fetal inflammatory response in preterm infants with chorioamnionitis and funisitis. *Pediatr Res.* 57:263-9.
Gruenheid and Finlay (2003) Microbial pathogenesis and cytoskeletal function. *Nature*, 422: 775-781.
Romero and Mazor (1988) Infection and preterm labor. *Clin Obstet Gynecol*; 31: 553-584.

Keywords: Placenta, chorioamnionitis, tight junction, occludin.