

## Identification of nuclear substrates of Akt/PKB by functional proteomics: prohibitin 2 is a target of Akt phosphorylation in human promyelocytic leukemia cells

Alberto Bavelloni<sup>1,2</sup>, Irene Faenza<sup>3</sup>, Manuela Piazzini<sup>3</sup>, William Blalock<sup>3</sup>, Antonietta D'Angelo<sup>1,2</sup>, Francesca Tagliavini<sup>3</sup>, Diego Pinetti<sup>4</sup>, Sandro Matteucci<sup>3</sup>, Giulia Adalgisa Mariani<sup>3</sup>, Lucio Cocco<sup>3</sup>

<sup>1</sup> RAMSES, Rizzoli Orthopedic Institute, Bologna, Italy

<sup>2</sup> Laboratory of Musculoskeletal Cell Biology, Rizzoli Orthopedic Institute, Bologna, Italy.

<sup>3</sup> Cellular Signaling Laboratory, Department of Human Anatomical Sciences, University of Bologna

<sup>4</sup> CIGS, University of Modena e Reggio Emilia, Italy

The serine/threonine protein kinase Akt is a major signal transducer of the phosphoinositide 3-kinase (PI 3-K) pathway in all cells and tissues and plays a pivotal role in the maintenance of cellular processes including cell growth, proliferation, survival, metabolism and development of many malignancies including acute myeloid leukemia. The frequent aberrant activation of the PI 3-K/Akt pathway in human cancer has made it an attractive therapeutic target. Therefore, the study of effector proteins downstream of Akt could clarify the role of Akt in the development of myeloid leukemia. Although both localization and activity of Akt in the nuclear compartment are well documented, most Akt substrates identified so far are located in the cytoplasm, while nuclear substrates have remained elusive. In this study, we applied a proteomic approach to identify novel Akt substrates by using an antibody that recognized a consensus motif phosphorylated by Akt (K/RXK/RXXS/T) when phosphorylated on S/T (anti-phospho-Akt substrate antibody). NB4 cells were treated with ATRA, and the putative Akt substrate proteins were isolated by immunoprecipitation with the anti-phospho-Akt substrate antibody. The proteins were separated on SDS-PAGE and analyzed by ESI-Q-TOF mass spectrometry.

This analysis indicated prohibitin 2, a potential tumor suppressor protein with potent transcriptional functions in the nucleus, as a putative substrate of Akt in the nucleus of NB4 cells. The putative Akt-Prohibitin 2 interaction was validated by reverse in vivo immunoprecipitation from nuclear protein of NB4 cells. In vitro phosphorylation of endogenous prohibitin 2 by recombinant Akt further validated this result.

Vandermoere F, El Yazidi-Belkoura I, Demont Y, Slomianny C, Antol J, Lemoine J, Hondemarck H. Proteomics exploration reveals that actin is a signaling target of the kinase Akt. *Mol Cell Proteomics*. 2007 Jan;6(1):114-24.

Kasashima K, Ohta E, Kagawa Y, Endo H. Mitochondrial functions and estrogen receptor-dependent nuclear translocation of pleiotropic human prohibitin 2. *J Biol Chem*. 2006 Nov 24;281(47):36401-10.

Keywords: topography, nucleus, Akt, leukemia, functional proteomics