

Safarpour Kapourchali, S., Maleki, M., Alizadeh Aliabadi, A., Rajaei, S., Faghihi, M. M. & Nasr Esfahani, M. (2025). Microbiota dynamic communities in sweet orange infected by “huanglongbing” in Iran. *Phytopathologia Mediter-ranea* 64(1): 129–143. doi: 10.36253/phyto-15662

Supplementary Table 1. Characteristics of geographical and climatic factors related to the sampling regions in Southern Iran.

Province	Geographical and Climatic factor					
	Latitude	Longitude	Elevation	Average temperatures (minimum/maximum)	Average annual precipitation	Average relative humidity
Fars (Darab)	28°75' N	54° 54' E	1131 m	11.5 to 36.5°C	220 mm	12–47% (38%)
Hormozgan (Rudan)	28° 52' N	57° 27' E	193 m	8 to 42°C	250 mm	17–67% (53%)
Kerman (Jiroft)	28° 67' N	57°74' E	650 m	8 to 38°C	250 mm	12–60% (42%)
Khuzestan (Dezful)	32°15' N	48° 24' E	150 m	7 to 43°C	372 mm	15–60% (50%)
Sistan and Baluchistan (Sarbaz)	26° 63' N	61° 25' E	868 m	2.5 to 43.5°C	227 mm	44–68% (55%)