

## RESEARCH PAPERS

# Bioassays on the activity of resveratrol, pterostilbene and phosphorous acid towards fungi associated with esca of grapevine

MAZZULLO ALESSANDRO<sup>1</sup>, STEFANO DI MARCO<sup>1</sup>, FABIO OSTI<sup>2</sup> and AUGUSTO CESARI<sup>2</sup>

<sup>1</sup>CNR, Centro di Studio dei Fitofarmaci, Via Filippo Re 8, 40126 Bologna, Italy

<sup>2</sup>Dipartimento di Protezione e Valorizzazione Agroalimentare, Università degli Studi, Via Filippo Re 8, 40126 Bologna, Italy

**Summary.** An examination was carried out under laboratory conditions to determine whether resveratrol, pterostilbene and mixtures of each stilbene with phosphorous acid had an inhibitory activity against some fungi involved in esca: *Phaeoacremonium aleophilum*, *Phaeoconiella chlamydospora*, *Libertella blepharis* (anamorph of *Eutypa lata*), *Fomitiporia punctata* and *Stereum hirsutum*. An inhibition of mycelial growth by resveratrol was not found but there was a clear correlation between pterostilbene concentration and reduction of radial growth confirming the antifungal activity of this phytoalexin. Although phosphorous acid proved to be generally ineffective, an increase of antifungal activity of stilbenes in the mixtures was observed with all fungi except *F. punctata*. The possible relation between fungal phenol oxidase activity and inhibition by mixtures of stilbenes and phosphorous acid was discussed.

**Key words:** grapevine, stilbenes, phosphorous acid, esca fungi, phenol oxidase activity.

## Introduction

Esca is a very destructive grapevine disease that is widespread in all grape-growing areas. Several micro-organisms are involved in a complex infection process not yet completely elucidated. The wood is colonized by different fungi acting in succession or in combination and producing various types of discoloration and decay (Mugnai *et al.*,

1996; Larignon and Dubos, 1997; Graniti *et al.*, 2000).

The discoloration normally precedes or occurs on the edge of decayed areas. Non-basidiomycetous fungi are consistently isolated from these discoloured regions, while white rot basidiomycetes are commonly found in zones with visible decay. Discoloration seems a necessary first stage before decay, and the mechanism by which such presumed preconditioning occurs is most commonly suggested to be via the detoxification of phenolic substances produced by the host as an infection response (Rayner and Boddy, 1988).

---

Corresponding author: S. Di Marco  
Fax: + 39 051 2091364  
E-mail: sdimarco@agrsci.unibo.it

The following pathogens are frequently isolated from necrosis typical of esca: *Phaeoacremonium chlamydosporum* (W. Gams, Crous, M.J. Wingf. & L. Mugnai), recently proposed by Crous and W. Gams (2000) as *Phaeomoniella chlamydospora* (W. Gams, Crous, M.J. Wingf. & L. Mugnai) Crous & W. Gams, *Phaeoacremonium aleophilum* (W. Gams, Crous, M.J. Wingf. & L. Mugnai), *Eutypa lata* (Pers.: Fr.) Tul. & C. Tul., *Fomitiporia punctata* (Murril) and *Stereum hirsutum* (Willd.: Fr.) S.F. Gray.

Stilbenes have long been considered important as conferring wood resistance to fungal decay: wood containing stilbenes frequently decomposes more slowly when exposed to various rot fungi (Hart and Shrimpton, 1979). Resveratrol is the major stilbene component in lignified stem tissue of grapevine and this compound probably also acts as a biosynthetic precursor of other *Vitis* stilbenes such as the viniferins and pterostilbene (Langcake and Pryce, 1977). *Trans*-pterostilbene in particular was reported by Langcake *et al.* (1979) to be a methylated resveratrol derivative with an interesting antifungal activity.

Under favourable infection conditions, esca-diseased vines treated with fosetyl Al compounds by wood injection showed positive results with a significant reduction in the severity of foliar symptoms. Although the treatment did not cure the vines, a positive long-term effect was that basic production criteria (weight of harvested bunches and Brix values) were preserved (Mazzullo *et al.*, 2000).

The action of fosetyl-Al in stimulating the host defence response as well as in having a direct inhibitory effect on the pathogens through phosphorous acid - to which fosetyl Al is rapidly degraded - was extensively described by Fenn and Coffey (1984).

A proper understanding of the relations between fosetyl-Al, the grapevine defence mechanism and esca pathogens appears to be of basic importance when evaluating of the above mentioned positive preliminary field results, if the potential of fosetyl-Al to control esca is to be correctly interpreted.

The aim of the present study was to determine the effect of resveratrol and pterostilbene on the mycelial growth of the fungi involved in the esca syndrome and to explore the role of these com-

pounds in mixtures with phosphorous acid. Phenol oxidase enzyme tests were also performed.

## Materials and methods

### Fungi and culture conditions

Cultures of *P. aleophilum*, *P. chlamydospora*, *Libertella blepharis* (anamorph of *Eutypa lata*), *F. punctata* and *Stereum hirsutum* isolated from esca-diseased plants were grown and maintained in the dark at 25±1°C on Petri plates containing potato dextrose agar (PDA, Difco, Detroit, MI, USA) at 39 g l<sup>-1</sup>. An isolate of each fungus was submitted to bioassays.

### Bioassays

Bioassays were carried out with resveratrol (3,4',5 - trihydroxy-trans-stilbene, 99%, Sigma, St Louis, MO, USA), pterostilbene (laboratory synthesis) and phosphorous acid (H<sub>3</sub>PO<sub>3</sub>, 99%, Aldrich, Milwaukee, WI, USA).

Pterostilbene (*trans*-3-5-dimethoxy-4'-hydroxystilbene) was produced in the laboratory by condensing *p*-hydroxybenzaldehyde with dimethoxy-3,5 phenylacetic acid in the presence of piperidine, in accordance with Massarani (1957). The last step of the synthesis involved a Perkin condensation which brought about a *trans* isomere with a coupling constant of 16.5 Hz between the vinyl protons (Pezet and Pont, 1988). The *trans* configuration of the pterostilbene was proved by NMR and mass spectrometry.

Resveratrol and pterostilbene were dissolved in absolute ethanol and the alcoholic solutions were added to a PDA substrate at 40°C. The final concentration of ethanol was 2% (v:v). An identical volume of ethanol was incorporated with the growth medium of the controls (Hart and Hillis, 1974; Bailey *et al.*, 1976; Hart and Shrimpton, 1979; Di Marco *et al.*, 1999).

Preliminary tests on the stilbenes were carried out with the following amounts of active ingredient in the medium: resveratrol at 10, 100, 500 and 1000 µg ml<sup>-1</sup>; pterostilbene at 1, 10 and 100 µg ml<sup>-1</sup>. At a later stage, for the study, phosphorous acid (300 µg ml<sup>-1</sup>), resveratrol (300 µg ml<sup>-1</sup>), resveratrol (300 µg ml<sup>-1</sup>) + phosphorous acid (300 µg ml<sup>-1</sup>), pterostilbene (10 µg ml<sup>-1</sup>), pterostilbene (10 µg ml<sup>-1</sup>) + phosphorous acid (300 µg ml<sup>-1</sup>) were dissolved in the medium. Resveratrol at concentrations above 100 ppm formed crystals.

The pH of the growth medium was measured by litmus paper (Farmitalia, Carlo Erba, Milano, Italy).

Inoculum plugs cut with a cork borer from the margin of growing cultures were placed at the centre of Petri plates containing treated medium. Control plugs were placed in untreated medium. Plates were incubated at 25±1°C in the dark. For each fungus colony diameters were measured at two points on each plate and averaged.

Treatments were replicated three times, each replicate consisting of one Petri-plate.

For each assessment date, data were analysed statistically using Duncan's multiple range test,  $P=0.05$ .

#### **Detection of resveratrol in growth medium**

The growth medium of plates treated with resveratrol or resveratrol + phosphorous acid and inoculated with *P. aleophilum* was analysed by Thin Layer Chromatography (TLC) to detect the presence of resveratrol. This analysis was performed because a brownish halo appeared around the edge of *P. aleophilum* colony grown on medium with resveratrol alone (i.e. without phosphorous acid).

Discs of PDA 4 mm in diameter were collected from plates containing medium + resveratrol and medium + resveratrol + phosphorous acid. Discs were cut either adjacent to the outer edge of the colony or away from the front of the mycelium. Samples were extracted in ethanol and analysed by TLC on silica gel plates (Baker-Flex IB2, Phillipsburg, NJ, USA) with cyclohexane/ethyl acetate (9:1) as the usual solvent system. Developed plates were detected by UV lamp (254 nm). Five discs per treatment were set up.

#### **Enzyme assays**

For each fungus the tests described below were performed according to Harkin and Obst (1973) and Rayner and Boddy (1988).

##### *Laccase*

I. A dilute 0.1% solution of syringaldazine (4-hydroxy-3,5-dimethoxybenzaldehyde azine, 99%, Aldrich) in absolute ethanol was added to PDA plates in which discs 5 mm in diameter from the margin of colonies of the pathogens were successively placed.

II. A dilute 0.1% solution of syringaldazine in absolute ethanol was placed dropwise with a capil-

lary pipette at the front of actively growing mycelium of pathogen colonies.

The laccase test reaction was shown by the medium turning deep purple (test I) or by the speed of the colour change (test II) due to syringaldazine oxidation by the laccase.

##### *Peroxidase*

The syringaldazine test was similar to laccase test II above. If no colour developed and laccase was absent, a drop of dilute hydrogen peroxide solution was placed on the edge of the colony. Peroxidase activity was shown by the appearance of a purple colour as with the laccase test.

##### *Tyrosinase*

A drop of 0.1% *p*-cresol (1.08 g) in 96% ethanol (100 ml) was added to the edge of the colony. Tyrosinase activity was shown by the appearance of an orange-brown coloration in the medium 4-48 hours after application.

##### *Catalase*

Catalase activity was carried out by placing a drop of hydrogen peroxide solution on the fungal cultures, and visually assessing the resulting effervescence.

The level of enzyme activity was expressed as "low" or "strong".

## **Results**

### **Bioassays**

#### *Preliminary assays of stilbenes*

Effects on mycelial growth by different amounts of resveratrol and pterostilbene active ingredient are summarized in Tables 1 and 2 respectively.

Resveratrol crystals formed when stilbene was used at 500 and 1000 µg ml<sup>-1</sup>. Crystals were uniformly dispersed in the medium and increased in number with increasing resveratrol concentration.

The morphological and growth characteristics of the colonies did not vary with the amount of resveratrol in the medium, whether as crystals or in diluted form.

Amounts of resveratrol up to 100 µg ml<sup>-1</sup> significantly reduced mycelial growth of *S. hirsutum* and, to a lesser extent, of *F. punctata*, compared to the control. No effects were observed with *L. blepharis*, *P. chlamydospora* or *P. aleophilum*. A

Table 1. Mycelial growth<sup>a</sup> of fungi with different amounts of resveratrol in the medium. Mycelial growth (mm) is expressed as colony diameter.

Fungal species	Resveratrol amount ( $\mu\text{g ml}^{-1}$ )	Days after treatment						
		2	5	8	11	14	17	20
<i>P. chlamydospora</i>	10	7.0 b <sup>b</sup>	11.3 b	14.7 b	19.3 a	25.3 bc	33.3 b	38.7 b
	100	11.0 c	16.0 c	20.3 c	25.7 b	29.0 c	36.0 b	41.0 b
	500	5.0 a	7.3 a	11.0 a	17.0 a	21.7 ab	25.0 a	28.3 a
	1000	5.0 a	7.3 a	10.3 a	15.0 a	17.3 a	21.0 a	25.3 a
	Control	5.0 a	10.7 b	13.3 ab	16.7 a	20.3 a	25.3 a	28.7 a
<i>P. aleophilum</i>	10	5.3 b	9.7 a	13.7 a	17.3 a	23.3 ab	29.7 bc	36.3 c
	100	7.0 c	10.7 a	15.0 a	19.3 a	27.0 b	33.3 c	40.3 c
	500	4.0 a	10.0 a	13.0 a	16.3 a	20.3 ab	25.7 ab	30.0 ab
	1000	4.0 a	8.3 a	10.7 a	14.7 a	17.7 a	21.7 a	26.0 a
	Control	4.0 a	9.0 a	12.0 a	17.0 a	22.3 ab	29.0 bc	35.3 bc
<i>L. blepharis</i>	10	7.7 a	16.7 a	48.0 b	70.3 a	-	-	-
	100	7.7 a	15.3 a	40.0 ab	63.3 a	-	-	-
	500	7.7 a	13.7 a	35.0 a	59.7 a	-	-	-
	1000	8.0 a	13.7 a	32.3 a	56.3 a	-	-	-
	Control	8.3 a	18.0 a	41.0 ab	71.7 a	-	-	-
<i>F. punctata</i>	10	4.0 a	13.0 bc	26.3 b	42.7 b	58.3 b	70.7 c	-
	100	4.0 a	15.0 c	29.7 c	45.3 b	63.0 b	72.7 c	-
	500	4.0 a	10.0 a	21.7 a	34.7 a	53.7 a	67.0 b	-
	1000	4.0 a	11.0 ab	23.0 a	35.0 a	52.0 a	62.7 a	-
	Control	4.0 a	12.7 bc	26.7 b	43.0 b	57.7 b	71.3 c	-
<i>S. hirsutum</i>	10	18.7 c	67.7 c	-	-	-	-	-
	100	16.0 b	62.7 bc	-	-	-	-	-
	500	14.3 ab	55.7 a	-	-	-	-	-
	1000	13.0 a	56.3 ab	-	-	-	-	-
	Control	18.0 c	62.3 bc	-	-	-	-	-

<sup>a</sup> Including the diameter (4 mm) of the inoculum plug.

<sup>b</sup> For each fungus, data in columns followed by the same letter do not differ significantly according to Duncan's MRT ( $P = 0.05$ ).  
-, Not recorded because colony had reached the edge of the plate.

statistically significant increase in growth rate was noted at 100  $\mu\text{g ml}^{-1}$  resveratrol for *P. chlamydospora* and, to a lesser extent, for *P. aleophilum* from the 14th day of inspection. In plates containing resveratrol, a brownish halo (pigmentation of the medium) appeared around the outer edge of *P. aleophilum* colony at 10 days from inoculation.

Pterostilbene inhibited mycelial growth of all fungi. On the whole, the amount of inhibition increased with the amount of pterostilbene in the medium. With *L. blepharis* this inhibition was statistically significant at all concentrations. With *S. hirsutum* and *F. punctata* inhibition was sig-

nificant at 10 and 100  $\mu\text{g ml}^{-1}$ . At the highest stilbene concentration, growth of *P. chlamydospora* was significantly lower than the control, with about 50% reduction. Pterostilbene was reasonably effective in reducing the growth of *P. aleophilum*; this reduction was statistically significant at concentration of 100  $\mu\text{g ml}^{-1}$ .

#### *Resveratrol + phosphorous acid*

Growth medium containing resveratrol formed crystals also when phosphorous acid was added.

Addition of phosphorous acid caused a lowering of the pH of the growth medium from 5.5 to 5.

Table 2. Mycelial growth<sup>a</sup> of fungi with different amounts of pterostilbene in the medium. Mycelial growth (mm) is expressed as colony diameter.

Fungal species	Pterostilbene amount ( $\mu\text{g ml}^{-1}$ )	Days after treatment						
		2	5	8	11	14	17	20
<i>P. chlamydospora</i>	1	4.0 a <sup>b</sup>	11.3 bc	14.0 bc	19.7 b	23.3 bc	25.7 b	28.7 b
	10	4.0 a	8.7 ab	12.0 b	17.0 b	21.7 b	22.0 b	27.0 b
	100	4.0 a	7.7 a	9.0 a	10.0 a	11.3 a	13.7 a	18.7 a
	Control	4.0 a	12.0 c	16.0 c	20.7 b	24.7 c	27.0 b	33.0 b
<i>P. aleophilum</i>	1	4.0 a	8.7 ab	10.7 b	13.3 b	18.3 b	21.7 a	29.0 bc
	10	4.0 a	7.3 a	10.3 ab	12.7 ab	17.7 b	21.7 a	26.3 ab
	100	4.0 a	7.3 a	9.0 a	11.3 a	14.7 a	20.3 a	25.3 a
	Control	4.0 a	9.3 b	11.7 b	15.3 c	18.7 b	24.3 b	31.0 c
<i>L. blepharis</i>	1	9.0 ab	34.7 c	58.7 c	78.3 c	-	-	-
	10	9.0 ab	25.3 b	42.0 b	66.0 b	-	-	-
	100	8.3 a	13.3 a	18.3 a	26.3 a	-	-	-
	Control	10.7 b	39.0 d	67.0 d	89.0 d	-	-	-
<i>F. punctata</i>	1	4.0 a	14.3 a	26.7 b	40.7 b	51.7 b	62.7 b	-
	10	4.0 a	12.7 a	23.3 a	35.0 a	45.3 a	54.7 a	-
	100	4.0 a	13.0 a	23.7 ab	34.3 a	44.0 a	53.7 a	-
	Control	4.0 a	12.7 a	26.0 ab	39.3 b	50.7 b	62.7 b	-
<i>S. hirsutum</i>	1	12.3 b	58.7 bc	-	-	-	-	-
	10	9.3 a	52.0 b	-	-	-	-	-
	100	7.7 a	20.0 a	-	-	-	-	-
	Control	14.0 b	62.0 c	-	-	-	-	-

a, b, - See Table 1.

Normally this mixture gave a significant reduction of mycelial growth with all fungi. With *F. punctata*, generally low inhibition only occasionally reached statistical significance (Table 3). The superior inhibition with the mixture compared to the compounds when tested separately, was very evident with *L. blepharis* and *P. aleophilum*, and to a lesser extent with *P. chlamydospora*. The brownish halo on a *P. aleophilum* medium was never observed when the medium was treated with resveratrol + phosphorous acid. Resveratrol crystals were observed at all inspection dates. *P. aleophilum* and *F. punctata* growth was stimulated to various degrees by phosphorous acid.

#### *Pterostilbene + phosphorous acid*

The pterostilbene + phosphorous acid mixture gave significant inhibition against all fungi at each

inspection dates. Growth of *L. blepharis*, *S. hirsutum* and, to a lesser extent, *P. aleophilum* was significantly lower than with the other treatments, at all inspection dates (Table 4). For *P. chlamydospora* growth inhibition with this mixture was significant from the sixth day of inspection, and *F. punctata* inhibition was also significant but less so. Phosphorous acid inhibited *L. blepharis*, *P. chlamydospora* and *P. aleophilum* but on the contrary stimulated *F. punctata*, *S. hirsutum* and *P. aleophilum* growth.

#### Detection of resveratrol in the growth medium

Resveratrol presence in the medium inoculated with *P. aleophilum* and treated with resveratrol was compared with that treated with resveratrol + phosphorous acid to account for the brownish halo appearance.

Table 3. Effect of resveratrol and phosphorous acid on fungal mycelial growth<sup>a</sup>. Mycelial growth (mm) is expressed as colony diameter.

Fungal species	Treatment	Amount of a.i. (µg ml <sup>-1</sup> )	Days after treatment																
			2	4	6	8	10	12	14	16	18	20							
<i>P. chlamydospora</i>	Resveratrol	300	5.0 a <sup>b</sup>	6.5 a	8.7 ab	12.0 a	16.0 a	19.3 ab	22.3 a	24.6 b	26.3 b	31.3 b							
	Phosphorous acid	300	5.0 a	7.3 a	11.0 b	13.7 a	15.7 a	18.3 a	20.3 a	24.0 b	25.7 b	27.3 ab							
	Resv. + phosph. ac.	300+300	5.0 a	5.7 a	7.0 a	10.0 a	11.7 a	14.0 a	17.0 a	18.0 a	20.7 a	22.3 a							
	Control	-	5.3 a	13.7 b	16.7 c	19.7 b	22.7 b	26.0 b	30.3 b	33.0 c	36.0 c	40.7 c							
<i>P. aleophilum</i>	Resveratrol	300	4.0 a	7.3 b	9.7 b	12.0 ab	14.3 ab	18.0 ab	21.0 b	22.7 ab	28.7 b	30.0 ab							
	Phosphorous acid	300	4.0 a	6.7 b	12.3 c	15.0 b	17.0 b	19.0 ab	19.7 ab	22.7 ab	24.3 ab	26.7 ab							
	Resv. + phosph. ac.	300+300	4.0 a	5.0 a	6.7 a	9.0 a	10.3 a	13.3 a	14.7 a	16.7 a	17.7 a	19.3 a							
	Control	-	4.0 a	7.0 b	9.3 b	12.0 ab	15.7 ab	21.7 b	27.0 b	29.3 b	31.3 b	33.0 b							
<i>L. blepharis</i>	Resveratrol	300	7.3 b	14.0 c	26.0 c	35.0 c	48.3 b	66.6 c	-	-	-	-							
	Phosphorous acid	300	4.0 a	5.0 b	6.3 b	9.0 b	11.3 a	19.3 b	-	-	-	-							
	Resv. + phosph. ac.	300+300	4.0 a	4.0 a	4.0 a	4.0 a	5.3 a	9.3 a	-	-	-	-							
	Control	-	8.3 b	18.7 d	28.0 c	42.0 d	54.7 b	81.7 d	-	-	-	-							
<i>F. punctata</i>	Resveratrol	300	5.7 ab	10.0 a	17.3 a	26.0 a	34.0 a	45.7 ab	53.7 a	64.0 a	69.3 b	-							
	Phosphorous acid	300	5.3 a	15.0 a	23.3 c	36.3 b	46.7 b	56.7 c	71.7 c	75.0 b	77.7 c	-							
	Resv. + phosph. ac.	300+300	5.3 a	12.0 a	17.3 a	26.3 a	33.3 a	43.0 a	52.0 a	60.3 a	66.3 a	-							
	Control	-	6.7 b	11.7 a	19.3 b	28.0 a	36.0 a	46.7 b	58.0 b	64.3 a	70.3 b	-							
<i>S. hirsutum</i>	Resveratrol	300	18.7 a	47.7 b	78.3 b	-	-	-	-	-	-	-							
	Phosphorous acid	300	23.0 b	56.7 c	82.7 b	-	-	-	-	-	-	-							
	Resv. + phosph. ac.	300+300	17.3 a	39.3 a	64.0 a	-	-	-	-	-	-	-							
	Control	-	22.0 b	56.7 c	83.3 b	-	-	-	-	-	-	-							

<sup>a, b</sup> - See Table 1.

Table 4. Effect of pterostilbene and phosphorous acid on fungal mycelial growth<sup>a</sup>. Mycelial growth (mm) is expressed as colony diameter (mm).

Fungal species	Treatment	Amount of a.i. ( $\mu\text{g ml}^{-1}$ )	Days after treatment																
			2	4	6	8	10	12	14	16	18	20							
<i>P. chlamydospora</i>	Pterostilbene	10	4.0 b <sup>b</sup>	7.3 a	9.3 a	12.3 ab	14.3 ab	18.7 ab	21.3 ab	22.0 ab	25.0 ab	27.3 ab							
	Phosphorous acid	300	4.7 b	7.0 a	11.0 ab	13.0 ab	15.0b c	17.7 ab	20.0 ab	23.0 bc	25.7 ab	26.6 ab							
	Ptero. + phosph. ac.	10+300	4.0 a	6.0 a	8.3 a	10.3 a	12.3 a	14.0 a	16.0 a	18.0 a	20.7 a	24.0 a							
	Control	-	6.0 b	8.7 a	13.7 b	17.0 b	18.3 c	23.7 b	26.0 b	28.3 c	31.0 b	34.3 b							
<i>P. aleophilum</i>	Pterostilbene	10	4.0 a	7.0 b	8.3 b	10.7 b	12.3 b	13.3 b	18.0 b	19.3 b	23.7 b	27.0 b							
	Phosphorous acid	300	4.0 a	6.7 b	12.3 c	15.0 c	17.0 c	19.0 c	19.7 b	22.7 b	24.3 b	26.7 b							
	Ptero. + phosph. ac.	10+300	4.0 a	4.0 a	5.0 a	6.0 a	7.7 a	8.7 a	10.0 a	10.7 a	13.3 a	14.7 a							
	Control	-	4.0 a	6.3 b	8.3 b	11.0 b	13.0 bc	14.7 b	18.3 b	22.3 b	26.7 b	33.7 b							
<i>L. blepharis</i>	Pterostilbene	10	6.0 b	14.7 c	24.3 c	31.3 c	50.3 c	68.0 c	-	-	-	-							
	Phosphorous acid	300	4.0 a	5.0 b	6.3 b	9.0 b	11.3 b	19.3 b	-	-	-	-							
	Ptero. + phosph. ac.	10+300	4.0 a	4.0 a	4.0 a	4.7 a	5.3 a	7.7 a	-	-	-	-							
	Control	-	8.3 c	24.0 d	44.3 d	56.3 d	71.3 d	88.7 d	-	-	-	-							
<i>F. punctata</i>	Pterostilbene	10	7.7 ab	14.0 a	24.3 a	33.0 ab	40.3 b	49.3 b	54.7 a	65.7 a	71.7 a	78.0 a							
	Phosphorous acid	300	7.3 ab	15.0 ab	23.3 a	35.3 c	46.7 c	56.7 c	71.7 c	75.0 b	77.7 b	82.7 b							
	Ptero. + phosph. ac.	10+300	7.0 a	13.7 a	23.3 a	31.7 a	38.7 a	47.7 a	54.0 a	65.7 a	72.3 a	76.3 a							
	Control	-	8.0 b	16.3 b	26.0 b	34.3 b	41.7 b	50.3 b	58.0 b	67.0 a	72.3 a	76.7 a							
<i>S. hirsutum</i>	Pterostilbene	10	11.3 a	35.3 b	70.0 b	-	-	-	-	-	-	-							
	Phosphorous acid	300	23.0 b	56.7 c	84.0 c	-	-	-	-	-	-	-							
	Ptero. + phosph. ac.	10+300	7.3 a	16.7 a	39.3 a	-	-	-	-	-	-	-							
	Control	-	22.0 b	51.0 c	85.7 c	-	-	-	-	-	-	-							

<sup>a, b</sup> - See Table 1.

Table 5. Phenol oxidase tests.

Fungal species	Laccase	Peroxidase	Tyrosinase	Catalase
<i>P. chlamydospora</i>	Negative	Negative	Negative	Negative
<i>P. aleophilum</i>	Negative	Negative	Negative	Strong
<i>L. blepharis</i>	Negative	Negative	Negative	Low
<i>F. punctata</i>	Strong	- <sup>a</sup>	Negative	-
<i>S. hirsutum</i>	Negative	Negative	Negative	Low

<sup>a</sup> Not determined.

In the pigmented area of the plates treated with resveratrol no crystals were visible and TLC analysis showed no stilbene was present; conversely, in the non pigmented area, the stilbene was present as crystals and was recovered by TLC analysis. In the medium treated with resveratrol + phosphorous acid there never was a brownish halo, and TLC showed stilbene was present.

#### Enzyme assays

Both laccase tests showed clearly that *F. punctata* was the only fungus that produced extracellular laccase. Neither peroxidase nor tyrosinase activity were detected with any of the fungi. *P. aleophilum* exhibited strong catalase activity as shown by immediate effervescence when dilute hydrogen peroxide was placed on culture. Catalase activity was weak for *L. blepharis* and *S. hirsutum*; completely absent for *P. chlamydospora* (Table 5).

#### Discussion

The aim of the study was to investigate the activity of resveratrol, pterostilbene and these same compounds + phosphorous acid towards fungi involved in esca.

A clear inhibition of mycelial growth by resveratrol was not found. Normally, the *in vitro* behaviour of this stilbene depended on the amount of it in the growth medium: at low levels (particularly 100 µg ml<sup>-1</sup>) it actually had a certain stimulant effect on mycelial growth of *P. chlamydospora* and *P. aleophilum* but at higher levels it inhibited growth, especially with *F. punctata* and *S. hirsutum*. These findings agree with the way in which

wood extractives interact with fungal activity either as stimulants or inhibitors of growth (Rayner and Boddy, 1988).

Pterostilbene was described by Langkake *et al.* (1979) as a *V. vinifera* phytoalexin and its antifungal activity was relatively high in comparison with resveratrol. Our data showed a direct correlation between the concentration of pterostilbene in the media and the inhibition of fungal radial growth.

A stilbene + phosphorous acid mixture was generally more inhibitory towards all the fungi except *F. punctata*. According to Mugnai *et al.* (1997) the insensitivity of this white-rot agent to the stilbenes is probably linked with its strong laccase activity. Besides, the involvement of oxidases in wood attack mechanism is well known: stilbenes in the hardwood of conifers are metabolized only by white-rot fungi producing laccase (Loman, 1970a, b). Phosphorous acid did not impair *F. punctata* growth in the presence of stilbenes.

The greater activity of the mixtures against the other fungi might be due to the lowering of the pH. Probably, resveratrol is taken up more readily by fungal cells when such lowering of the pH occurs (Deverall and Rogers, 1972). Moreover, Hoos and Blaich (1990) established that the strongest activity of resveratrol against *Botrytis* on solid medium was at pH 5.

Despite the lack of laccase, peroxidase and tyrosinase activity, *P. aleophilum* was able to inactivate the resveratrol. Although the way in which resveratrol disappeared in the brownish halo is still unclear, the oxidation of this stilbene in the brown area was probably related to the strong catalase activity of this fungus. The superior activity of resveratrol + phosphorous acid appeared to be closely linked to the inability of *P. aleophilum* to metabolize this stilbene in the presence of phosphorous acid.

A clear inhibitory effect of phosphorous acid was not found with any fungus except *L. blepharis*. Phosphorous acid did not act as an antifungal agent as such but enhanced stilbene activity.

Although the relation between the *Vitis* stilbenes (resveratrol and pterostilbene) and resistance to fungal infections is still unclear, the results obtained in this study demonstrated that a mixture stilbenes + phosphorous acid was more active against *L. blepharis*, *P. aleophilum*, *P. chlamydospora* and *S. hirsutum*. Basically, the combina-

tion of phosphorous acid and the stilbenes may contribute to prevent or reduce fungal infection. Further and more extensive *in vivo* investigations are obviously needed to evaluate the potential of fosetyl-Al, to control the esca syndrome, especially considering the spread of young grapevine decline due to *Phaeoacremonium* spp. (Ferreira *et al.*, 1994; Morton, 1995, 1997; Bertelli *et al.*, 1998; Pascoe, 1998; Scheck *et al.*, 1998).

## Literature cited

- Bailey J.A., G.A. Carter and R.A. Skipp, 1976. The use and interpretation of bioassays for fungitoxicity of phytoalexins in agar media. *Physiological Plant Pathology*, 8, 189–194.
- Bertelli E., L. Mugnai and G. Surico, 1998. Presence of *Phaeoacremonium chlamydosporum* in apparently healthy rooted grapevine cuttings. *Phytopathologia Mediterranea*, 37, 79–82.
- Crous P.W. and W. Gams, 2000. *Phaeomoniella chlamydospora* gen. et comb. nov., a causal organism of Petri grapevine decline and esca. *Phytopathologia Mediterranea*, 39(1), 112–118.
- Deverall B.J. and P.M. Rogers, 1972. The effect of Ph and composition of test solutions on the inhibitory activity of wyerone acid towards germination of fungal spores. *Annals of Applied Biology*, 72, 301–305.
- Di Marco S., A. Mazzullo, F. Calzarano and A. Cesari, 1999. *In vitro* studies on the phosphorous acid-vitis stilbenes interaction, and *in vivo* phosetyl Al activity towards *Phaeoacremonium* spp. grapevine wood decay agents. In: Modern Fungicides and Antifungal Compounds II (H. Lyr, P.E. Russel, H.W. Dehene, H.D. Sisler eds). Intercept Ltd, Hampshire, UK, 171–177.
- Fenn M.E. and M.D. Coffey, 1984. Studies on the *in vitro* and *in vivo* antifungal activity of fosetyl-Al and phosphorous acid. *Phytopathology*, 74(5), 606–611.
- Ferreira J.H.S., P.S van Wyk. and E. Venter 1994. Slow dieback of grapevine: association of *Phialophora parasitica* with slow dieback of grapevines. *South African Journal of Enology and Viticulture*, 15(1), 9–11.
- Graniti A., G. Surico and L. Mugnai, 2000. Esca of grapevine: a disease complex or a complex of diseases? *Phytopathologia Mediterranea*, 39(1), 16–20.
- Harkin J.M. and J.R. Obst, 1973. Syringaldazine, an effective reagent for detecting laccase and peroxidase in fungi. *Experientia*, 29(4), 381–387.
- Hart J.H. and W.E. Hillis, 1974. Inhibition of wood rotting fungi by stilbenes and other polyphenols in *Eucalyptus sideroxyton*. *Phytopathology*, 64, 939–948.
- Hart J.H. and D.M. Shrimpton, 1979. Role of stilbenes in resistance of wood to decay. *Phytopathology*, 69, 1138–1143.
- Hoos G. and R. Blaich, 1990. Influence of resveratrol on germination of conidia and mycelial growth of *Botrytis cinerea* and *Phomopsis viticola*. *Journal of Phytopathology*, 129, 102–110.
- Langkake P. and R.J. Price, 1977. A new class of phytoalexins from grapevines. *Experientia*, 33, 151–152.
- Langkake P., C.A. Cornford and R.J. Price, 1977. Identification of pterostilbene as a phytoalexin from *Vitis vinifera* leaves. *Phytochemistry*, 18, 1025–1027.
- Larignon P. and B. Dubos, 1997. Fungi associated with esca disease in grapevine. *European Journal of Plant Pathology*, 103, 147–157.
- Loman A.A., 1970a. The effect of heartwood fungi of *Pinus contorta* var. *latifolia* on pinosylvin, pinosylvinmonomethyl ether, pinobanksin and pinocembrin. *Canadian Journal of Botany*, 48, 737–747.
- Loman A.A., 1970b. Bioassays of fungi isolated from *Pinus contorta* var. *latifolia* with pinosylvin, pinosylvinmonomethyl ether, pinobanksin and pinocembrin. *Canadian Journal of Botany*, 48, 1303–1308.
- Massarani E., 1957. Ulteriori ricerche sopra derivati stilbenici e difeniletanici. *Il Farmaco*, 12(5), 380–386.
- Mazzullo A., F. Calzarano, S. Di Marco and A. Cesari, 2000. Experiences on the control of esca disease fungi by fosetyl Al. In: Integrated Control in Viticulture. *Bullettin IOBC/WPRS*, 23(4), 77–78.
- Morton L., 1995. Mystery disease hits young vines. *Wines and vines* 76(11), 46–47.
- Morton L., 1997. Update on black goo. *Wines and vines*, 78(1), 62–64.
- Mugnai L., A. Graniti and G. Surico, 1999. Esca (black measles) and brown wood-streaking: two old and elusive diseases of grapevines. *Plant Disease*, 83(5), 404–418.
- Mugnai L., G. Surico and A. Sfalanga, 1997. Produzione di enzimi esocellulari da parte di funghi del legno di viti colpita da “mal dell’esca”. *Micologia italiana*, 1, 11–22.
- Pascoe I., 1998. Trunk disease of grapevines – perspectives from a tour of California. *Australian Grapegrower Winemaker*, 417, 68–71.
- Pezet R. and V. Pont, 1988. Mise en évidence de ptérostilbène dans les grappes de *Vitis vinifera*. *Plant Physiology and Biochemistry*, 26(5), 603–607.
- Rayner A.D.M. and L. Boddy, 1988. Fungal decomposition of wood. Its biology and ecology. John Wiley & Sons Ltd., Bath, UK, 587 pp.
- Scheck H., S. Vasquez, D. Fogle and W.D. Gubler, 1998. Grape growers report losses to black foot and grapevine decline. *California Agriculture*, 52(4), 19–23.

Accepted for publication: 26 June, 2000