

## Viruses infecting figs in Egypt

ESAM KAMAL FAHMY ELBESHEHY<sup>1</sup> and TOUFIC ELBEAINO<sup>2</sup>

<sup>1</sup>Agricultural Botany Department, Faculty of Agriculture, Suez Canal University, 41522 Ismailia, Egypt

<sup>2</sup>Istituto Agronomico Mediterraneo di Bari, Via Ceglie 9, 70010 Valenzano (Bari), Italy

**Summary.** Fig production in Egypt is in continuous decline because many diseases, particularly those caused by viruses, are compromising this crop. RT-PCR assays were conducted on 60 fig samples collected from three Egyptian fig-growing provinces (Ismailia, Qena and North Sinai), from the three main fig varieties (cvs. Sultany, Abode and El-Adasy), to investigate the presence of *Fig leaf mottle-associated virus 1* (FLMaV-1), *Fig leaf mottle-associated virus 2* (FLMaV-2), *Fig mild mottle-associated virus* (FMMaV) and *Fig mosaic virus* (FMV). The overall average incidence of infection reached 90%, with a peak of 95% in Ismailia province. All four viruses were detected in tested samples, with infection rates of 68.3% for FLMaV-1, 35% for FLMaV-2, 28.3% for FMMaV and 46.7% for FMV. This is the first report of the presence of these viruses in Egypt and offers a preliminary insight on the unsatisfactory sanitary status of fig in this country.

**Key words:** *Ficus carica*, FMV, virus detection, RT-PCR, electron microscopy.

### Introduction

According to FAO statistics, Egypt is the second largest fig producer in the world, after Turkey, with a production of 170,000 tons (Anonymous, 2005). More than 5 million fig trees are grown in Egypt on a total area of 32,060 ha (General Administration for Agriculture, Statistics, MOA, Egypt). In the last few years, poor growth of fig trees, scant yield and low fruit quality have been common complaints from growers in different regions of Egypt. The unknown sanitary status of fig crops in Egypt, together with the traditional techniques for fig propagation (grafting and/or self-rooted cuttings), have favored the spread of pathogens, especially viruses. Symptoms of mosaic disease, always presumed and only recently confirmed to be of viral origin (Elbeaino *et al.*, 2009a), are scattered in gardens as well as in commercial fig or-

chards throughout the country. The first description of fig mosaic disease (MD) was from California by Condit and Horn (1933), and this disease was found to be transmitted by grafting and in nature by the eryophid mite *Aceria ficus* (Flock and Wallace, 1955). The aetiology of this disease remained uncertain for a long time, even though isometric and filamentous virus particles were occasionally observed in thin-sectioned tissues of symptomatic leaves of fig accessions from Italy and England (Martelli *et al.*, 1993). Putative potyviruses were also recorded from Croatia (Grbelja and Eric, 1983) and Spain (Serrano *et al.*, 2004), a putative carlavirus from Japan (Doi, 1989), and double stranded RNAs (dsRNA) with sizes ranging from 0.6 to 6.6 kb were recovered from infected trees in Portugal and Turkey (Nolasco and Sequeira, 1991; Açıkğös and Döken, 2003).

A closterovirus denoted *Fig leaf mottle-associated virus 1* (FLMaV-1) was the first virus infecting fig in nature whose partial nucleotide sequence was determined and submitted in Genbank (Elbeaino *et al.*, 2006). Later records of

---

Corresponding author: T. Elbeaino  
Fax: +39 080 4606275  
E-mail: elbeaino@iamb.it

fig-infecting viruses rapidly increased and additional viruses infecting fig joined the list: *Fig leaf mottle-associated virus 2* (FLMaV-2), *Fig mosaic virus* (FMV), *Fig latent virus 1* (FLV-1), *Fig mild mottle-associated virus* (FMMaV) and *Fig cryptic virus* (FCV) (Elbeaino *et al.*, 2007; 2009a; 2010; 2011; Gattoni *et al.*, 2009). Contemporarily, other viruses likely belonging to *Partitiviridae* (Luteovirus-like) and *Caulimoviridae* (Badnavirus-like) families, with scanty molecular information (short nucleotide sequences) of their genomes, were also found in infected diseased figs (Walia *et al.*, 2009, Tzanetakakis *et al.*, 2009). For all of these viruses, molecular information on genome organization and sequences which permit their detection using molecular tools are now available. In this paper we report a preliminary investigation carried out in some fig-growing areas of southern Egypt to assess the presence of FLMaV-1, FLMaV-2, FMMaV and FMV using molecular tools (RT-PCR).

## Materials and methods

### Field survey and plant material

A total of sixty samples was collected from naturally infected fig trees of cvs. Sultany, Abode

and El-Adasy in the three fig-growing provinces of Egypt, Ismailia, Qena and North Sinai. Sampling was conducted during May 2010, and collections were made both from symptomless and MD-symptomatic trees. Symptoms varied from chlorotic blotches, vein clearing, vein banding, chlorosis, mosaic and chlorotic ringspot, all suggestive of virus-like diseases (Figure 1). All of the samples were assayed by RT-PCR to assess the presence of FLMaV-1, FLMaV-2, FMMaV and FMV in Egypt.

### Total nucleic acid extraction (TNA)

Total nucleic acids (TNAs) were extracted from 100 mg of leaf veins or cortical scrapings by ground each sample in 1 mL of grinding buffer (4.0M guanidine thiocyanate, 0.2M NaOAc pH 5.2, 25mM EDTA, 1.0M KOAc and 2.5% (w/v) PVP-40), and were silica-purified (Foissac *et al.*, 2001).

### cDNA synthesis

Eight to 10  $\mu\text{L}$  of TNA extracts were mixed with 1  $\mu\text{L}$  of random hexamer primers, (Boehringer Mannheim GbmH, Mannheim, Germany) ( $0.5 \mu\text{g } \mu\text{L}^{-1}$ ), denatured at  $95^\circ\text{C}$  for 5 min and quickly chilled on ice. Reverse-transcription was done for 1 h at  $39^\circ\text{C}$  by adding 4  $\mu\text{L}$  M-MLV buffer 5 $\times$  (50 mM Tris-HCl pH 8.3, 75 mM KCl, 3 mM  $\text{MgCl}_2$ ), 2  $\mu\text{L}$  of 10mM DTT, 0.5  $\mu\text{L}$  of 10 mM dNTPs, and




Figure 1. Symptoms of fig-mosaic diseased plants observed in inspected fig orchards and showing a wide range of foliar discoloration and malformation symptoms. (a) vein feathering, (b) vein banding, (c) leaf deformation, (d) chlorosis, (e) mosaic, (f) chlorotic blotching, (g) chlorotic ringspot and (h) leaf puckering.

200 units of *Moloney Murine Leukaemia virus* (M-MLV) reverse transcriptase (Bethesda Research Laboratories, Gaithersburg, M, USA) in a final volume of 20  $\mu$ L.

### PCR

The detection of FLMaV-1, FLMaV-2, FMMaV and FMV in RT-PCR was conducted using four sets of specific primers, whose nucleotide sequences and use-conditions were previously described (Elbeaino *et al.*, 2006, 2007, 2009a, 2010). Briefly, for each sample, 2.5  $\mu$ L of reverse-transcribed TNA mixture was submitted to amplification with the addition of 2.5  $\mu$ L of 10 $\times$  Taq polymerase buffer (Promega Corporation, Madison, WI, USA), with a final concentration of 1.5 mM MgCl<sub>2</sub> for a total volume of 25  $\mu$ L. PCR products were analyzed by electrophoresis on 1.2% agarose gel prepared in 1 $\times$  TBE buffer (Sambrook *et al.*, 1989), stained with

ethidium bromide and examined on a UV transilluminator.

### Electron microscope

For thin sectioning, tissue pieces from veins and mesophyll tissues of the discolored areas of young leaves were processed according to standard procedures (Martelli and Russo, 1984), i.e. fixation in 4% glutaraldehyde in 0.05M phosphate buffer for 2 h, post-fixation in 1% osmium tetroxide for 2 h, staining overnight in 2% aqueous uranyl acetate, dehydration in ethanol, and embedding in Spurr's medium. Thin sections were stained with lead citrate and viewed with a JOEL-JEA100 CX electron microscope Unit (Faculty of Science, Zagazig University, Alsharqiya, Egypt). Controls consisted of leaf tissues from a PCR-negative fig seedling processed as above.




Figure 2. Electropherogram of polyacrylamide gel showing PCR amplifications from infected figs. (A) FLMaV-1, (B) FLMaV-2, (C) FMMaV and (D) FMV. Lane M, DNA ladder marker, Lanes 1, 2 and 3 represent PCR-positive infected fig plants. Lanes 4 and 5 represent healthy figs.

Table 1. Incidence of *FLMaV-1*, *FLMaV-2*, *FMMaV* and FMV infections in three fig-growing provinces of Egypt as determined by RT-PCR assays.

Province	Tested trees No.	Infected trees		<i>FLMaV-1</i>		<i>FLMaV-2</i>		<i>FMMaV</i>		FMV	
		No.	%	No.	%	No.	%	No.	%	No.	%
Ismailia	20	19	95	18	90	5	25	3	15	9	45
Qena	20	17	85	9	45	9	45	7	35	10	50
Sinai	20	18	90	14	70	7	35	7	35	9	45
Total	60	54	90	41		21		17		28	
Mean infection rate			90		68.3		35		28.3		46.7

## Results and discussion

### RT-PCR Detection

RT-PCR assays of samples yielded four DNA amplifications of expected sizes, 352 bp, 360 bp, 311 bp and 300 bp, as results for *FLMaV-1*, *FLMaV-2*, *FMMaV* and FMV infections, respectively (Figure 2). From a total of 60 samples, 54 (90%) were infected by at least one virus. *FLMaV-1* was the prevailing virus with an infection rate of 68.3% (Table 1). The incidence of this virus was particularly high on cv. Sultany grown in Ismailia province (90%) and on cv. El-Adasy in Sinai (70%). FMV ranked second as incidence (46.7%) and was substantially equally distributed in all cultivars and regions (45–50%). *FLMaV-2* and *FMMaV*, the other two viruses checked in this

study, although to a lesser extent were found in all three provinces, with peaks of infection of 45% for *FLMaV-2* in Qena province and of 35% for *FMMaV* in Qena and Sinai provinces. These results are in harmony with those previously reported in other Mediterranean countries (Elbeaino *et al.*, 2006; 2007; 2009b; 2009c). Unlike closterovirus infections, which were frequently detected in asymptomatic trees, all FMV PCR-positive samples were closely correlated with mosaic symptoms in diseased fig plants.

### Electron microscope

Electron micrographs of sectioned cells revealed the presence of double membrane bodies (DMBs), considered to be FMV particles




Figure 3. Groups of Double-membraned Bodies (DMB) in the cytoplasm of mesophyll cells from a naturally infected fig. Arrows indicate the aggregates of convoluted electron-dense double membrane elements. Ch, chloroplast, CW, cell wall, Bar = 500 nm.

with dimensions of 50–80 nm (Figure 3). Such structures were not found in PCR-negative samples to FMV, thus confirming previous reports on the aetiology of DMB in mosaic affected figs (Elbeaino *et al.*, 2009b). These structures are a consistent feature of diseased figs, regardless of the variety and the country of origin (Plavsic and Milicic 1980; Martelli *et al.*, 1993; Appiano *et al.*, 1995; Castellano *et al.*, 2007; Elbeaino *et al.*, 2009a, 2009b).

The outcome of this preliminary work extends the knowledge on the spread of fig viruses in the Mediterranean region, particularly in Egypt for which no information was previously available. This is the first report of FLMaV-1, FLMaV-2, FMMAV and FMV occurring in the Egyptian fig orchards. Although this assessment was limited to 60 trees, the results obtained clearly indicate how the sanitary status of fig crop has deteriorated in Egypt (90% of viral infection). Particularly worrying is the incidence of FMV, since this has proved to be the unique virus closely correlated with the FMD (Elbeaino *et al.*, 2009a). High incidence of FMV is not surprising considering the way this virus spreads in figs through infected propagating material (cuttings and grafting), and natural vectors (eriphyid mites). In Egypt there is no information on the presence of *Aceria ficus* (Eriophyidae) and *Planococcus ficus* (Pseudococcidae), the recognized vectors of FMV and FLMaV-2, respectively. However, such presence in the Egyptian orchards would likely aggravate the sanitary status and the level of infections in the surveyed areas.

The several FMV-infected samples found in association with most of the mosaic symptoms in the field further confirms what was previously reported regarding the aetiology of FMV. Nevertheless, this postulate was infringed by a few cases where FMV was detected in symptomless fig trees. Whether this is due to the virus strain present in the country or to the biological response of some Egyptian fig varieties to FMV infection remains to be determined.

The knowledge we have gained on the incidence of virus diseases of fig in Egypt provides information on which a sanitary selection, sanitation and certification programs can be initiated for the production of healthy propagating plant material of fig in this country.

## Literature cited

- Açikgös S. and M.T. Döken, 2003. The determination of sampling time for dsRNA isolation of the agent of fig mosaic disease prevalent in Aegean region-Turkey. *Acta Horticulturae* 605, 307–310.
- Anonymous, 2005. Food and Agriculture Organization of United Nations, Statistical Data (FAOSTAT) (Available online at <http://faostat.fao.org/>.)
- Appiano A.M., M. Conti and N. Zini, 1995. Cytopathological study of the double-membrane bodies occurring in fig plants affected by fig mosaic disease. *Acta Horticulturae* 386, 585–592.
- Castellano M.A., G. Gattoni, A. Minafra, M. Conti and G.P. Martelli, 2007. Fig mosaic in Mexico and South Africa. *Journal of Plant Pathology* 89, 441–443.
- Condit I.J. and W.T. Horne, 1933. A mosaic of the fig in California. *Phytopathology* 23, 887–896.
- Doi Y., 1989. Fig virus S. In: *Directory and Dictionary of Animal, Bacterial and Plant Viruses*, (R. Hull, F. Brown, C. Payne, ed.). Nature Publishing Group, London, UK, pp. 76.
- Elbeaino T., M. Digiario, A. De Stradis and G.P. Martelli, 2006. Partial characterization of a closterovirus associated with a chlorotic mottling of fig. *Journal of Plant Pathology* 88, 187–192.
- Elbeaino T., M. Digiario, A. De Stradis and G.P. Martelli, 2007. Identification of a second member of the family Closteroviridae in mosaic-diseased figs. *Journal of Plant Pathology* 89, 119–124.
- Elbeaino T., M. Digiario, A. Alabdullah, A. De Stradis, A. Minafra, N. Mielke, M.A. Castellano and G.P. Martelli, 2009a. A multipartite single-stranded negative-sense RNA virus is the putative agent of fig mosaic disease. *Journal of General Virology* 90, 1281–1288.
- Elbeaino T., M. Digiario and G.P. Martelli, 2009b. Complete nucleotides sequence of four viral RNAs segments of fig mosaic virus. *Archives of Virology* 154(11), 1719–1727.
- Elbeaino T., S. Nahdi, M. Digiario, A. Alabdullah and G.P. Martelli, 2009c. Detection of Fig leaf mottle-associated virus 1 and Fig leaf mottle-associated virus 2 in the Mediterranean region and study on sequence variation of the hsp70 gene. *Journal of Plant Pathology* 91, 425–431.
- Elbeaino T., K. Heinoun, M. Digiario and G.P. Martelli, 2010. Fig mild mottle-associated virus, a novel closterovirus infecting fig. *Journal of Plant Pathology* 92, 165–172.
- Elbeaino T., R. Abou Kubaa, M. Digiario, A. Minafra and G.P. Martelli, 2011. The complete nucleotide sequence and genome organisation of fig cryptic virus, a novel bipartite dsRNA virus infecting fig, widely distributed in the Mediterranean basin. *Virus Genes* 42(3), 415–421.
- Flock R.A. and J.M. Wallace, 1955. Transmission of fig mosaic by the eriophyid mite *Aceria ficus*. *Phytopathology* 45, 52–54.
- Foissac X., L. Svanella-Dumas, P. Gentit, M.J. Dulucq and T. Candresse, 2001. Polyvalent detection of fruit tree Tricho, Capillo and Foveavirus by nested RT-PCR us-

- ing degenerated and inosine containing primers (DOP RT-PCR). *Acta Horticulturae* 550, 37–43.
- Gattoni G., A. Minafra, M.A. Castellano, A. De Stradis, D. Boscia, T. Elbeaino, M. Digiario and G.P. Martelli, 2009. Some properties of Fig latent virus 1, a new member of the family Flexiviridae. *Journal of Plant Pathology* 91, 543–552.
- Grbelja J. and Z. Eric, 1983. Isolation of a potyvirus from *Ficus carica* L. *Acta Botanica Croatia* 42, 11–14.
- Martelli G.P., M.A. Castellano and R. Laforteza, 1993. An ultrastructural study of fig mosaic. *Phytopathologia Mediterranea* 32, 33–43.
- Martelli G.P. and M. Russo, 1984. Use of thin sectioning for visualization and identification of plant viruses. *Methods in Virology* 8, 143–224.
- Nolasco G. And O.A. de Sequeira, 1991. Double-stranded RNA (dsRNA) associated with fig mosaic disease. In: *Proceedings 4th Portuguese-Spanish Biochemistry Congress*, Lisbon, Portugal, 5 P2-Mo.
- Plavšić B. and D. Milicic, 1980. Intracellular changes in trees infected with fig mosaic. *Acta Horticulturae* 110, 281–286.
- Sambrook J., E.F. Fritsch and T. Maniatis, 1989. *Molecular cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA.
- Serrano, L., J. Ramon, J. Segarra, V. Medina, M.A. Achón, M. López and M. Juárez, 2004. New approach in the identification of the causal agent of fig mosaic disease. *Acta Horticulturae* 657, 559–566.
- Tzanetakakis I. and R. Martin, 2009. New viruses identified in fig trees exhibiting fig mosaic disease. *Proceedings 21st International Conference on Virus and Other Graft Transmissible Diseases of Fruit Crops*, 5–10 July, 2009, Neustadt/Weinstrasse, Germany, 29.
- Walia J.J., N.M. Salem and W.B. Falk, 2009. Partial sequence and survey analysis identify a multipartite, negative-sense RNA virus associated with fig mosaic. *Plant Disease* 93, 4–10.

*Accepted for publication: April 5, 2011*