

RESEARCH PAPERS - 10TH SPECIAL ISSUE ON GRAPEVINE TRUNK DISEASES

Pathogenicity of ten *Phaeoacremonium* species associated with esca and Petri disease of grapevine

MAHLATSE A. BALOYI^{1,2}, LIZEL MOSTERT² and FRANCOIS HALLEEN^{1,2}

¹ Plant Protection Division, ARC Infruitec-Nietvoorbij, Private Bag X5026, Stellenbosch, 7599, South Africa

² Department of Plant Pathology, University of Stellenbosch, Private Bag X1 Matieland, 7602, South Africa

Summary. Nineteen species of *Phaeoacremonium* have been associated with grapevines in South Africa, of which only six species have been confirmed as pathogens through pathogenicity tests conducted on field-grown grapevines. This study determined the pathogenic status of ten *Phaeoacremonium* spp. recently found for the first time on South African grapevines. These were: *Pm. australiense*, *Pm. austroafricanum*, *Pm. fraxinopennsylvanicum*, *Pm. griseo-olivaceum*, *Pm. griseorubrum*, *Pm. iranianum*, *Pm. italicum*, *Pm. prunicolum*, *Pm. scolyyti* and *Pm. sicilianum*. In the pathogenicity tests, *Ph. parasiticum* was used as the positive control, and sterile water as the negative control. Up to three isolates were used per species, depending on isolate availability. Freshly cut pruning wounds in a 9-year-old Cabernet Sauvignon vineyard in Stellenbosch, South Africa, were inoculated with 200 conidia of each fungus per wound. Inoculated pruning wounds were removed after 18 months, cut longitudinally and lesion lengths were measured. Re-isolation proportions were determined by conducting isolations from inoculated spurs. All the inoculated isolates successfully colonized pruning wounds, and caused lesions that were significantly different from the negative control. All isolates were re-isolated at proportions varying from 28.6 to 85.7%. *Phaeoacremonium griseo-olivaceum* STE-U 7859 produced the longest lesions (mean = 79.5 mm) and *Pm. iranianum* STE-U 6998 the shortest (62.0 mm). No statistically significant differences in mean lesion lengths were observed between the inoculated species. There were also no significant differences between isolates of the same species, except in *Pm. prunicolum* where isolate STE-U 5968 produced longer lesions (mean = 77.3 mm) than STE-U 7857 (62.3 mm). This study confirmed the capabilities of all the tested *Phaeoacremonium* spp. to infect grapevine pruning wounds and cause lesions. The study also confirmed the importance of pruning wounds as ports of entry by these pathogens into host plants.

Keywords: *Vitis* spp., grapevine trunk diseases, pruning wound infections.

Introduction

Of the 61 *Phaeoacremonium* W. Gams, Crous & M.J. Wingf. species known, 33 have been isolated from grapevines (Gramaje *et al.*, 2015; Spies *et al.*, 2018). Species of *Phaeoacremonium* are known inhabitants of many woody hosts and have also been isolated from humans (Ajello *et al.*, 1974; Crous *et al.*, 1996; Mostert *et al.*, 2005; Damm *et al.*, 2008; Gramaje *et al.*, 2012; Marin-Felix *et al.*, 2019). Some of the species associated with grapevine have also been found from fruit and

nut trees, including economic important pome and stone fruit species, olives, kiwifruit, guava, fig, pomegranate, loquat, persimmon, mulberry, date palm and medlar, as well as numerous ornamental trees and shrubs, some occurring in forests, plantations, gardens or parks (Gramaje *et al.*, 2015; Spies *et al.*, 2018).

Phaeomoniella (*Pa.*) *chlamydospora* (W. Gams, Crous, M.J. Wingf. & Mugnai) Crous & W. Gams, together with *Phaeoacremonium* (*Pm.*) spp., cause Petri disease (Scheck *et al.*, 1998; Mugnai *et al.*, 1999), which is commonly found in young grapevines of approx. 1-5 years age (Halleen and Groenewald, 2005). Species of *Phaeoacremonium* are well adapted endophytes, capable of becoming pathogenic when vines are sub-

Corresponding author: F. Halleen
E-mail: halleenf@arc.agric.za

jected to stress (Scheck *et al.*, 1998; Ferreira *et al.*, 1999). External host decline symptoms include leaf necrosis, shoot dieback, shortened internodes or (in extreme cases) the vine death. Internal symptoms include brown streaks in xylem tissues, seen in a longitudinal cuts infected stems, or the black/brown gummoses exuded from transversally cut wounds (Scheck *et al.*, 1998; Ferreira *et al.*, 1999; Mugnai *et al.*, 1999). *Phaeoacremonium* spp. are also associated with esca of grapevines. Esca is caused by the Petri disease pathogens together with wood rotting fungi of the Hymenochaetales (Mugnai *et al.*, 1999; White *et al.*, 2011a; Cloete *et al.*, 2015a, b). The brown streaking found in Petri diseased and esca-affected grapevines is due to the accumulation of phenolic compounds and tyloses in infected tissues, which block the xylem vessels and impair translocation of water and nutrients (Ferreira *et al.*, 1999; Edwards *et al.*, 2007).

Petri disease pathogens infect grapevines primarily through susceptible pruning wounds (Larignon and Dubos, 2000; Eskalen *et al.*, 2007; Van Niekerk *et al.*, 2011). Sucker wounds have also been shown to be susceptible, and could act as additional infection sites during spring and early summer (Makatini, 2014). Fruiting bodies of the pathogens formed within vineyards release spores mostly during and after rainfall periods, although spore release can occur without rainfall (Edwards and Pascoe, 2001; Eskalen *et al.*, 2005a; b; Rooney-Latham *et al.*, 2005). Spore trapping studies conducted in Western Cape (South Africa) vineyards and rootstock mother plant nurseries have shown the presence of air-borne spores within these vineyards (Baloyi, 2016). Spore release also coincided with winter and spring pruning periods, as well as the period when rootstock canes were harvested in nurseries. Studies in France, California and Italy have also showed the availability of air-borne spores during pruning periods (Larignon and Dubos, 2000; Eskalen and Gubler, 2001; Quaglia *et al.*, 2009). Winter pruning activities in the Western Cape occur between June and August, whereas de-suckering activities (winter pruning), the removal of unwanted shoots from vine trunks, cordons and spurs, occurs in spring and early summer (September to November). Rootstock canes are harvested between April and July. Multiple wounds made on each vine during these periods increase the risk of vines becoming infected. Other modes of pathogen spread have recently been reported, including arthropods which vector Petri disease pathogens to pruning wounds in vineyards (Moyo *et al.*,

2014), infected pruning shears under greenhouse conditions (Agustí-Brisach *et al.*, 2015), and the use of infected planting material (Fourie and Halleen, 2002; Halleen *et al.*, 2003; Fourie and Halleen, 2004).

In California, pruning wound susceptibility was shown to last for up to 16 weeks, which would enable even slow-growing pathogens to colonize and infect the wounds before they heal (Eskalen *et al.*, 2007). In France, wounds remained susceptible to *Pm. minimum* (Tul. & C. Tul.) Gramaje, L. Mostert & Crous for 7-9 weeks in the early dormant season, and for only 2 weeks in late winter pruning periods when bleeding occurred (Larignon and Dubos, 2000). Furthermore, Munkvold and Marois (1995) found that vines pruned towards the end of dormancy release exudates that contained carbohydrates, amino acids and organic acids, which promoted rapid growth of microflora that compete with pathogens. Other factors such as vine training method, as for bilateral cordons, result in multiple wounds and more pathogen entry sites, which may allow multiple infections (Gu *et al.*, 2005). Pruning spurs to two buds has been shown to promote rapid infection of cordons, as lesions extended quicker into the cordons and trunks from the inoculated pruning wounds (Eskalen *et al.*, 2007; Halleen *et al.*, 2007). To date, no grapevine cultivar has been shown to be resistant to *Phaeoacremonium* spp., or to any of the trunk disease pathogens. Pruning wound protection is, therefore, advocated as a disease management strategy (Halleen *et al.*, 2010; Kotze *et al.*, 2011; Mutawila *et al.*, 2016). However, the increasing number of fungal pathogens from unrelated fungal genera means that single products or modes of action may not be effective for control of the complex of grapevine trunk disease pathogens.

At commencement of this project, 17 *Phaeoacremonium* spp. were isolated from grapevines in South Africa. These were: *Pm. Minimum*; *Pm. italicum* Carlucci & M.L. Raimondo (reported as *Pm. alvesii*); *Pm. austroafricanum* L. Mostert, W. Gams & Crous; *Pm. iranianum* L. Mostert, Gräfenhan, W. Gams & Crous; *Pm. krajdinii* L. Mostert, Summerb. & Crous; *Pm. fraxinopennsylvanicum* (T.E. Hinds) D. Gramaje, L. Mostert & Crous; *Pm. parasiticum* (Ajello, Georg & C.J.K. Wang) W. Gams, Crous & M.J. Wingf.; *Pm. scolyti* L. Mostert, Summerb. & Crous; *Pm. sicilianum* Essakhi, Mugnai, Surico & Crous; *Pm. subulatum* L. Mostert, Summerb. & Crous; *Pm. venezuelense* L. Mostert, Summerb. & Crous; *Pm. viticola* J. Dupont.; *Pm. australiense* L. Mostert, Summerb. & Crous; *Pm. griseorubrum* L.

Mostert, Summerb. & Crous; *Pm. griseo-olivaceum* (Damm, L. Mostert & Crous) Gramaje, L. Mostert & Crous; *Pm. inflatipes* W. Gams, Crous & M.J. Wingf.; and *Pm. prunicolum* L. Mostert, Damm & Crous (Mostert *et al.*, 2005, 2006b; White *et al.*, 2011b; Spies *et al.*, 2018). Of these, only *Pm. minimum*, *Pm. krajdenii*, *Pm. parasiticum*, *Pm. subulatum*, *Pm. venezuelense* and *Pm. viticola* have been subjected to pathogenicity studies on grapevines in South Africa (Halleen *et al.*, 2007). The present study was therefore conducted to assess the pathogenic status of ten of these newly found *Phaeoacremonium* species, on grapevine pruning wounds under field conditions.

Materials and methods

Isolate selection and inoculum preparation

The *Phaeoacremonium* isolates used in this study were previously recovered and identified in spore trapping studies (Baloyi, 2016) and disease incidence surveys conducted in South African vineyards (Mostert *et al.*, 2005, 2006b; White *et al.*, 2011b) (Table 1). The species tested included *Pm. australiense*, *Pm. austroafricanum*, *Pm. italicum* (previously reported as *Pm. alvesii*), *Pm. fraxinopennsylvanicum*, *Pm. griseo-olivaceum*, *Pm. griseorubrum*, *Pm. iranianum*, *Pm. prunicolum*, *Pm. scolyti* and *Pm. sicilianum*. *Phaeoacremonium parasiticum* was used as the positive control (Halleen *et al.*, 2007) in the pathogenicity tests, and sterile water was used as the negative control. Up to three strains per isolate were used, depending on availability. Isolates were plated onto Potato Dextrose Agar with chloramphenicol (PDA-C), and grown at 25°C for 2 weeks. The cultures were then each flooded with 20 mL of sterile distilled double autoclaved water. Conidia were dislodged from the mycelia using a sterile glass rod, and the conidia suspensions were filtered with doubled cheese cloth. The conidia concentration was adjusted to 10⁴ mL⁻¹ after enumeration using a haemocytometer.

Pruning wound inoculations

The inoculation trial was conducted from August 2013 and March 2015, in a 9-year-old Cabernet Sauvignon vineyard at the Nietvoorbij Campus of ARC Infruitec-Nietvoorbij in Stellenbosch, Western Cape Province, South Africa. The trial was set up in a completely randomized block design, with 27 treatments

each replicated 15 times. Pruning wounds were the experimental unit. The vines were cordon trained and spur pruned to two buds. Five pruning wounds were made per vine and individual vines received one treatment. Due to sap flow, pruning wounds were not inoculated immediately, but within 24 h of pruning they were each inoculated with 20 µL of conidium suspension (200 conidia per wound).

Trial evaluation

The trial was evaluated after 18 months from the date of inoculation. Stubs with inoculated pruning wounds were removed, individually placed in plastic bags and immediately taken to the laboratory for assessment and to conduct isolations. The stubs were each cut longitudinally to measure lesion length with a caliper. The stubs were then surface sterilized by immersing into 70% ethanol for 30 s, followed by 1 min in 3.5% sodium hypochlorite solution and again for 30 s in 70% ethanol. Pieces of small tissue sections (1 × 1 × 2 mm) were each aseptically dissected with a sterile scalpel from just below the wound scar interface, as well as along the length of the lesion, and plated onto PDA-C plates. The inoculated Petri dishes were maintained at 25°C for 4 weeks and closely monitored for any *Phaeoacremonium* spp. growth. Re-isolated pathogens were identified based on morphological characteristics and then verified by randomly selecting two representative isolates from each treatment for sequencing. DNA was extracted with CTAB buffer according to Damm *et al.* (2008). The partial beta-tubulin gene regions were amplified with PCR using primers T1 and Bt2b (Glass and Donaldson, 1995; O' Donnell and Cigelink, 1997), as described by Mostert *et al.* (2006b). PCR products were cleaned using an MSB Spin PCRapase kit (Invitex), and the cleaned products were sequenced with the same primers using ABI PRISM Big Dye Terminator v3.1 Cycle Sequencing Ready Reaction Kit (PE Biosystems). The products were then analyzed on an ABI Prism 3130XL DNA sequencer (Perkin-Elmer). Sequences were compared to reference sequences of each species in the megablast function of the NCBI's GenBank nucleotide database (www.ncbi.nlm.nih.gov), to confirm identity. Mean lesion lengths were calculated, and subjected to one way analyses of variance (ANOVA) using SAS. Student's t-test for least significant difference (LSD) was calculated (at $P \leq 0.05$) to separate means.

Table 1. *Phaeoacremonium* spp. and isolates used in the pathogenicity study.

<i>Phaeoacremonium</i> spp.	Accession number	Host	Location of origin
<i>Pm. australiense</i>	STE-U 7863	Spore trap ^a	Slanghoek, Western Cape
	STE-U 7862	Spore trap ^a	Slanghoek
	STE-U 7861	Spore trap ^a	Slanghoek
<i>Pm. austroafricanum</i>	LM 733	<i>Vitis vinifera</i> ^b	Wellington, Western Cape
<i>Pm. italicum</i>	STE-U 6988	<i>Vitis vinifera</i> ^c	Klawer, Western Cape
	STE-U 6989	<i>Vitis vinifera</i> ^c	Klawer
	STE-U 7000	<i>Vitis vinifera</i> ^c	De Rust, Western Cape
<i>Pm. fraxinopennsylvanicum</i>	STE-U 6987	<i>Vitis vinifera</i> ^c	Hermanus, Western Cape
<i>Pm. griseorubrum</i>	STE-U 7881	<i>Vitis</i> sp.	Wellington
	STE-U 7882	<i>Vitis</i> sp.	Wellington
	STE-U 7856	<i>Vitis</i> sp.	Wellington
<i>Pm. griseo-olivaceum</i>	STE-U 7860	Spore trap ^a	Durbanville, Western Cape
	STE-U 7859	Spore trap ^a	Durbanville
	STE-U 7858	Spore trap ^a	Durbanville
<i>Pm. iranianum</i>	STE-U 6998	<i>Vitis vinifera</i> ^c	Calitzdorp, Western Cape
	STE-U 6999	<i>Vitis vinifera</i> ^c	Calitzdorp
<i>Pm. prunicolum</i>	STE-U 5968	<i>Prunus salicina</i> ^d	Mookgopong, Limpopo
	STE-U 7857	Spore trap ^a	Stellenbosch, Western Cape
<i>Pm. scolyti</i>	STE-U 7854	Spore trap ^a	Slanghoek
	STE-U 7855	Spore trap ^a	Rawsonville, Western Cape
	STE-U 7876	Spore trap ^a	Rawsonville
<i>Pm. sicilianum</i>	STE-U 7879	Spore trap ^a	Rawsonville
	STE-U 7880	Spore trap ^a	Rawsonville
	STE-U 7877	Spore trap ^a	Rawsonville
<i>Pm. parasiticum</i>	STE-U 7875	Spore trap ^a	Paarl, Western Cape
	STE-U 7878	Spore trap ^a	Stellenbosch

^a Isolates were obtained as aerial spore inoculum in vineyards (Baloyi, 2016).

^b Mostert *et al.* (2006b).

^c White *et al.* (2011b).

^d Damm *et al.* (2008).

Results

Mean lesion lengths of vascular discolouration caused by the 27 treatments, and mean re-isolation percentages, are presented in Table 2. All the isolates caused black to brown discolourations, as vascular streaking inside the inoculated spurs. All isolates

were considered pathogenic, as they produced lesion lengths significantly different to the negative experimental control ($P < 0.001$), and some lesions extended beyond the inoculated pruning wounds into the cordons. There was no significant variation in mean lesion lengths produced by each species. *Phaeocr-*

Table 2. Mean lesion lengths and re-isolation proportions for *Phaeoacremonium* spp. inoculated on Cabernet Sauvignon pruning wounds.

<i>Phaeoacremonium</i> spp.	Mean lesion length (mm)	Mean re-isolation (%)	Accession number
<i>Pm. griseo-olivaceum</i>	79.53 a	78.57	STE-U 7859
<i>Pm. prunicolum</i>	77.27 ab	64.29	STE-U 5968
<i>Pm. griseo-olivaceum</i>	75.54 abc	80.00	STE-U 7858
<i>Pm. parasiticum</i> (positive control)	74.53 abc	78.57	STE-U 7875
<i>Pm. sicilianum</i>	74.44 abc	76.92	STE-U 7880
<i>Pm. australiense</i>	74.30 abc	73.33	STE-U 7863
<i>Pm. griseorubrum</i>	73.27 abc	69.23	STE-U 7881
<i>Pm. australiense</i>	73.06 abc	42.86	STE-U 7861
<i>Pm. griseorubrum</i>	72.95 abc	69.23	STE-U 7882
<i>Pm. sicilianum</i>	72.04 abc	76.92	STE-U 7877
<i>Pm. scolyti</i>	71.59 abc	85.71	STE-U 7854
<i>Pm. scolyti</i>	71.51 abc	84.62	STE-U 7855
<i>Pm. sicilianum</i>	71.51 abc	80.00	STE-U 7879
<i>Pm. italicum</i>	70.91 abc	35.71	STE-U 7000
<i>Pm. iranianum</i>	70.05 abc	71.43	STE-U 6999
<i>Pm. austroafricanum</i>	67.97 abc	69.23	LM 733
<i>Pm. italicum</i>	67.54 abc	28.57	STE-U 6988
<i>Pm. griseo-olivaceum</i>	67.33 abc	84.62	STE-U 7860
<i>Pm. scolyti</i>	65.77 abc	66.67	STE-U 7876
<i>Pm. fraxinopennsylvanicum</i>	65.65 abc	78.57	STE-U 6987
<i>Pm. parasiticum</i> (positive control)	65.21 abc	57.14	STE-U 7878
<i>Pm. italicum</i>	64.50 bc	53.85	STE-U 6989
<i>Pm. australiense</i>	64.39 bc	57.14	STE-U 7862
<i>Pm. griseorubrum</i>	64.17 bc	73.33	STE-U 7856
<i>Pm. prunicolum</i>	62.26 c	57.14	STE-U 7857
<i>Pm. iranianum</i>	62.00 c	60.00	STE-U 6998
Negative control (sterile water)	14.46 d	0	
LSD ($P = 0.05$)	14.93		

^a Different letters indicate significant differences to the negative control ($P < 0.001$).

emonium griseo-olivaceum strain STE-U 7859 produced the longest lesions (mean = 79.5 mm). The shortest lesions (mean = 62.0 mm) were produced by *Pm. iranianum* strain STE-U 6998. Positive control wounds inoculated with two isolates of *Pm. parasiticum* pro-

duced mean lesion lengths of, respectively, 74.5 and 65.2 mm. All the inoculated isolates produced lesion lengths similar to *Pm. parasiticum*, a known pathogen, and therefore all the isolates are confirmed as pathogenic to grapevine. There were no significant differ-

ences between isolates of the same species, except for *Pm. prunicolum*, where STE-U 5968 produced significantly longer lesions (mean = 77.3 mm) than STE-U 7857 (62.3 mm).

All isolates were re-isolated from inoculated pruning wounds after 18 months. The isolates with the greatest proportions of re-isolation were STE-U 7854 and STE-U 7855 (*Pm. scolyti*), STE-U 7860 and STE-U 7858 (*Pm. griseo-olivaceum*) and STE-U 7879 (*Pm. sicilianum*), with re-isolation proportions of 89.0 to 84.6%. Two of the three *Pm. italicum* isolates were re-isolated at the least proportions of, respectively, 28.6 and 35.7% (Table 2).

Discussion

This study showed that all 11 inoculated *Phaeoacremonium* species were pathogenic and capable of causing vascular discolouration when inoculated onto grapevine pruning wounds. The ten *Phaeoacremonium* species with unknown pathogenicity to grapevines all formed lesions that were not significantly different in mean lengths from *Pm. parasiticum*, a known pathogen of grapevines. Of the species tested, *Pm. fraxinopennsylvanicum*, *Pm. iranianum*, *Pm. scolyti* and *Pm. sicilianum* have been tested on grapevines in other grapevine-producing countries (Gramaje *et al.*, 2007; Aroca and Raposo, 2009; Gramaje *et al.*, 2009; Gramaje *et al.*, 2010; Mohammadi and Banihashemi, 2012; Özben *et al.*, 2012; Úrbez-Torrez *et al.*, 2014; Mohammadi and Hashemi, 2015). All these species caused vascular discolourations inside inoculated grapevine shoots, which is consistent with the findings of the present study. Furthermore, reduced root weight, chlorotic leaves, severe defoliation and wilting symptoms have been observed from grapevine shoots inoculated with *Pm. fraxinopennsylvanicum* (Gramaje *et al.*, 2007; Aroca and Raposo, 2009; Úrbez-Torrez *et al.*, 2014). Low mean shoot weights were also reported in grapevine shoots inoculated with *Pm. iranianum* and *Pm. sicilianum* (Gramaje *et al.*, 2009).

All *Phaeoacremonium* spp. from South African grapevine, previously described as *Pm. aloesii* (White *et al.*, 2011b), were re-identified by Spies *et al.* (2018) as *Pm. italicum* based on the recent finding of this new species from Italian vineyards (Raimondo *et al.*, 2014). Although these isolations were made from declining and esca-affected vines in both countries, pathogenicity studies have not been previously conducted. The present report is, therefore, the first confirmation of

Pm. italicum as grapevine pathogen. *Phaeoacremonium austroafricanum* has only been reported from grapevines in South Africa (Mostert *et al.*, 2006b), but this is also the first confirmation of the pathogenic status of this species. Until the recent finding in South Africa (Spies *et al.*, 2018), *Phaeoacremonium griseo-olivaceum* and *Pm. prunicolum* have not been previously reported on grapevines. *Phaeoacremonium australiense* has previously only been isolated from grapevines in Australia and Uruguay (Mostert *et al.*, 2005; Abreo *et al.*, 2011), and *Pm. griseorubrum* from Italian vineyards (Essakhi *et al.*, 2008), but pathogenicity studies have not been conducted for these species. Pathogenicity has only been tested on *Prunus armeniaca* L. and *P. salicina* Lindl. in South Africa (Damm *et al.*, 2008). This is, therefore, the first confirmation of *Phaeoacremonium griseo-olivaceum*, *Pm. prunicolum*, *Pm. australiense* and *Pm. griseorubrum* as grapevine pathogens. These species were also trapped as airborne spores in Western Cape vineyards during the 2012 and 2013 seasons (Baloyi, 2016). The presence of these species as aerial inoculum, and confirmation of their capability to cause symptoms in grapevines and stone fruit, highlights the risks of establishing stone fruit orchards in close proximity to vineyards. This is also true for *Pm. iranianum* and *Pm. scolyti* (Damm *et al.*, 2008; Baloyi, 2016). This conclusion agrees with a study that showed *Pm. minimum* isolated from apple trees could infect grapevine shoots and develop lesions (Cloete *et al.*, 2011). Furthermore, *Phaeoacremonium minimum* isolates from apple, apricot and grapevine were pathogenic to apple, inducing wood discolouration on apple stems (Arzanlou *et al.*, 2014). Aroca and Raposo (2009) also used isolates of *Pm. inflatipes* from *Quercus* and *Pm. fraxinopennsylvanicum* from *Fraxinus* to show the capability of these species to infect and cause discolouration in grapevines. *Phaeoacremonium* spp. have wide host ranges, which favours the survival and spread of these pathogens, as several woody hosts can serve as pathogen reservoirs (Spies *et al.*, 2018). Cross dispersal between orchards and / or alternative hosts and vineyards is therefore of great concern, due to the increase in species diversity, and to the increasing risk of mating types being introduced that could result in production of sexual structures on vines.

All species of *Phaeoacremonium* tested on pruning wounds in this study were successfully re-isolated, although this was at varying proportions between species and isolates. The least re-isolation proportions were obtained with two of the three *Pm. italicum* iso-

lates. This species has previously only been isolated from grapevines in South Africa and Italy (White *et al.*, 2011b; Spies *et al.*, 2018). Spies *et al.* (2018) recently isolated *Pm. italicum* from eight additional hosts in South Africa, and Carlucci *et al.* (2015) also isolated this fungus from diseased olive trees in Italy. Infrequent release and low spore numbers in vineyards (Baloyi, 2016), isolation only from a limited number of grapevines (White *et al.*, 2011b), and presence in several other hosts may indicate that this species has been introduced more recently, or that grapevine is not its primary host.

There were statistically significant difference in virulence between the *Phaeoacremonium* isolates tested in this study, except for *Pm. prunicolum* where differences were observed between the two isolates. The most virulent isolate of *Pm. prunicolum* (STE-U 5968) was previously isolated from *Prunus salicina* in the Limpopo Province (Damm *et al.*, 2008), and was significantly different from the isolate found as airborne inoculum during spore trapping studies conducted in the Western Cape Province (Baloyi, 2016). The two isolates may be genetically different, resulting in different levels of virulence. Another explanation for the apparent lack of virulence differences in most of the *Phaeoacremonium* spp. may be that several lesions extended into the inoculated cordons, which could not be measured using the methods adopted for this study.

Isolates used in the present study were obtained from different geographic locations, and were only tested in one vineyard in Stellenbosch, which is, in some instances, environmentally different from the other regions from which isolates were collected. The capability of isolates from all these regions to infect grapevine pruning wounds, irrespective of their origins, and in the case of *Pm. prunicolum* also from another host, highlights the highly adaptive character of these pathogens. This shows the importance of movement of plant material between regions, which can introduce new pathogenic species into areas where they have not been previously reported. This is particularly the case for species such as *Pm. sicilianum* and *Pm. australiense*, which have not been reported from Stellenbosch vineyards (White *et al.*, 2011b; Baloyi, 2016) but have been shown to cause infections of grapevine pruning wounds in this region.

As was reported by Halleen *et al.* (2007), in the present study, many lesions extended from the inoculated two bud spurs into the cordons. This demonstrates

the risk associated with this pruning method, and how rapidly pruning wound infections can spread into the cordons. Other pruning practices, such as double pruning, have been previously been reported as a strategy to manage infection by *Eutypa lata* (Pers.) Tul. & C. Tul. (Weber *et al.*, 2007), but this practice has been widely used in South Africa for many years without any apparent reduction in the incidence of trunk diseases.

The pathogenicity trial in this study was not repeated. Pathogenicity of *Pm. fraxinopennsylvanicum*, *Pm. iranianum*, *Pm. scolyti* and *Pm. sicilianum* have been tested on grapevines in other countries, and the present study confirmed that South African isolates are pathogenic. Results obtained with the remaining six species, which are here confirmed as grapevine pathogens for the first time, were similar to previous pathogenicity studies conducted on grapevine pruning wounds in Stellenbosch (Halleen *et al.*, 2007).

This study confirms pruning wounds as entry sites for infection by *Pm. australiense*, *Pm. austroafricanum*, *Pm. fraxinopennsylvanicum*, *Pm. griseorubrum*, *Pm. griseo-olivaceum*, *Pm. iranianum*, *Pm. italicum*, *Pm. prunicolum*, *Pm. scolyti* and *Pm. sicilianum*, irrespective of the geographic origins of the isolates. The importance of pruning wound protection in vineyards and rootstock mother fields is thus emphasized, and this should form the foundation of an integrated management strategy to combat esca and Petri diseases of grapevine.

Acknowledgements

The authors acknowledge funding received from the South African Agricultural Research Council (ARC), National Research Foundation (NRF), Department of Science and Technology (DST), Technology and Human Resources for Industry Programme (THRIP) and Winetech (Project WW06/41). Technical assistance was provided by the ARC-Infruitec-Niet-voorbij technical team.

Literature cited

- Abreo E., S. Martinez and L. Bettucci, 2011. *Phaeoacremonium chlamydospora* and *Phaeoacremonium* spp. in grapevines from Uruguay. *Phytopathologia Mediterranea* 50, S77–S85.
- Agustí-Brisach C., M. Leon, J. García-Jiménez and J. Armengol, 2015. Detection of grapevine trunk pathogens on pruning shears and evaluating of their potential for spread of infection. *Plant Disease* 99, 976–981.

- Ajello L., L.K. Georg, R.T. Steigbigel and C.J.K. Wang, 1974. A case of phaeohyphomycosis caused by a new species of *Phialophora*. *Mycologia* 66, 490–498.
- Andolfi A., L. Mugnai, J. Luque, G. Surico, A. Cimmino and A. Evidente, 2011. Phytotoxins produced by fungi associated with grapevine trunk diseases. *Toxins* 3, 1569–1605.
- Aroca A. and R. Raposo, 2009. Pathogenicity of *Phaeoacremonium* species on grapevine. *Journal of Phytopathology* 157, 413–419.
- Arzanlou M., A. Narmani, S. Khodaei and S. Moshari, 2014. Pome and stone fruit trees as possible reservoir hosts for *Phaeoacremonium* spp., the causal agents of grapevine esca disease, in Iran. *Archives of Phytopathology and Plant Protection* 47, 717–727.
- Baloyi M.A., 2016. *Inoculum Ecology of Petri Disease Fungi in Grapevines of South Africa*. PhD dissertation, University of Stellenbosch, South Africa. 162 pp.
- Carlucci A., F. Lops, F. Cibelli and M.L. Raimondo, 2015. *Phaeoacremonium* species associated with olive wilt and decline in southern Italy. *European Journal of Plant Pathology* 141, 717–729.
- Cloete M., P.H. Fourie, U. Damm, P.W. Crous and L. Mostert, 2011. Fungi associated with die-back symptoms of apple and pear trees, a possible inoculum source of grapevine trunk disease pathogens. *Phytopathologia Mediterranea* 50, S175–S190.
- Cloete M., M. Fischer, L. Mostert and F. Halleen, 2015a. Hymenochaetales associated with esca-related wood rots on grapevine with a special emphasis on the status of esca in South African vineyards. *Phytopathologia Mediterranea* 54, 299–312.
- Cloete M., L. Mostert, M. Fischer and F. Halleen, 2015b. Pathogenicity of South African Hymenochaetales taxa isolated from esca-infected grapevines. *Phytopathologia Mediterranea* 54, 368–379.
- Crous P.W., W. Gams, M.J. Wingfield and P.S. Van Wyk, 1996. *Phaeoacremonium* gen. nov. associated with wilt and declining diseases of woody host and human infections. *Mycologia* 88, 786–796.
- Damm U., L. Mostert, P.W. Crous and P.H. Fourie, 2008. Novel *Phaeoacremonium* species associated with necrotic wood of *Prunus* trees. *Persoonia* 20, 87–102.
- Di Marco S., F. Calzarano, W. Gams and A. Cesari, 2000. A new wood decay of kiwifruit in Italy. *New Zealand Journal of Crop Horticulture Science* 28, 69–72.
- Edwards J. and I.G. Pascoe, 2001. Pycnidial state of *Phaeoconiella chlamydospora* found on Pinot Noir grapevines in the field. *Australasian Plant Pathology* 30, 67.
- Edwards J., I.G. Pascoe and S. Salib, 2007. Impairment of grapevine xylem function by *Phaeoconiella chlamydospora* infection is due to more than physical blockage of vessels with 'goo'. *Phytopathologia Mediterranea* 46, 87–90.
- Eskalen A. and W.D. Gubler, 2001. Association of spores of *Phaeoconiella chlamydospora*, *Phaeoacremonium inflatipes* and *Pm. aleophilum* with grapevine cordons in California. *Phytopathologia Mediterranea* 40, S429–S432.
- Eskalen A., S. Rooney-Latham and W.D. Gubler, 2005a. First report of perithecia of *Phaeoacremonium viticola* on grapevine and ash tree (*Fraxinus latifolia*) in California. *Plant Disease* 89, 686.
- Eskalen A., S. Rooney-Latham and W.D. Gubler, 2005b. Occurrence *Togninia fraxinopennsylvanica* on Esca-diseased grapevines (*Vitis vinifera*) and declining ash trees (*Fraxinus latifolia*) in California. *Plant Disease* 89, 528.
- Eskalen A., A.J. Feliciano and W.D. Gubler, 2007. Susceptibility of grapevine pruning wounds and symptom development in response to infection by *Phaeoacremonium aleophilum* and *Phaeoconiella chlamydospora*. *Plant Disease* 91, 1100–1104.
- Ferreira J.H.S., P.S. Van Wyk and F.J. Calitz, 1999. Slow dieback of grapevine in South Africa: Stress-related predisposition of young vines for infection by *Phaeoacremonium chlamydospora*. *South African Journal of Enology and Viticulture* 20, 43–46.
- Fourie P.H. and F. Halleen, 2002. Investigation on the occurrence of *Phaeoconiella chlamydospora* in canes of rootstock mother vines. *Australasian Plant Pathology* 31, 425–426.
- Fourie P.H. and F. Halleen, 2004. Occurrence of grapevine trunk disease pathogens in rootstock mother plants in South Africa. *Australasian Plant Pathology* 33, 313–315.
- Glass N.L. and G.C. Donaldson, 1995. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61, 1323–1330.
- Gramaje D., S. Alaniz, A. Pérez-Sierra, P. Abad-Campos, J. García-Jiménez and J. Armengol, 2007. First Report of *Phaeoacremonium mortoniae* causing Petri Disease of Grapevine in Spain. *Plant Disease* 91, 1206.
- Gramaje D., J. Armengol, M. Colino, R. Santiago, E. Moralejo, D. Olmo, J. Luque and L. Mostert, 2009. First report of *Phaeoacremonium inflatipes*, *P. iranianum* and *P. sicilianum* causing Petri disease of grapevine in Spain. *Plant Disease* 93, 964.
- Gramaje D., J. García-Jiménez and J. Armengol, 2010. Field evaluation of grapevine rootstocks inoculated with fungi associated with Petri disease and esca. *American Journal of Enology and Viticulture* 61, 512–520.
- Gramaje D., C. Agustí-Brisach, A. Pérez-Sierra, E. Moralejo, D. Olmo, L. Mostert, U. Damm and J. Armengol, 2012. Fungal trunk pathogens associated with wood decay of almond trees on Mallorca (Spain). *Persoonia* 28, 1–13.
- Gramaje D., L. Mostert, J.Z. Groenewald and P.W. Crous, 2015. *Phaeoacremonium*: from esca disease to phaeohyphomycosis. *Fungal Biology* 119, 759–783.
- Gu S., R.C. Cochran, G. Du, A. Hakim, K.C. Fugelsang, J. Ledbetter, C.A. Ingles and P.S. Verdegaal, 2005. Effect of training-pruning regimes on Eutypa dieback and performance of Cabernet Sauvignon grapevines. *Journal of Horticultural Science and Biotechnology* 80, 313–318.
- Halleen F. and M. Groenewald, 2005. Grapevine diagnostic observations with special reference to trunk diseases. In: *The 15th Biennial Australasian Plant Pathology Society Conference Handbook*. September 26–29, 2005, Geelong, Victoria, Australia, p.142 (abstract).
- Halleen F., P.W. Crous and O. Petrini, 2003. Fungi associated with healthy grapevine cuttings in nurseries, with special reference to pathogens involved in the decline of young vines. *Australasian Plant Pathology* 32, 47–52.
- Halleen F., L. Mostert and P.W. Crous, 2007. Pathogenicity testing of lesser-known vascular fungi of grapevines. *Australasian Plant Pathology* 36, 277–285.

- Halleen F., P.H. Fourie and P.J. Lombard, 2010. Protection of grapevine pruning wounds against *Eutypa lata* by biological and chemical methods. *South African Journal of Enology and Viticulture* 31, 125–132.
- Kotze C., J.M. Van Niekerk, L. Mostert, F. Halleen and P.H. Fourie, 2011. Evaluation of biocontrol agents for grapevine pruning wound protection against trunk pathogen infection. *Phytopathologia Mediterranea* 50, S247–S263.
- Larignon P. and B. Dubos, 2000. Preliminary studies on the biology of *Phaeoacremonium*. *Phytopathologia Mediterranea* 39, 184–189.
- Lynch S.C., P.J. Zambino, J.S. Mayorquin, D.H. Wang and A. Eskalen, 2013. Identification of new fungal pathogens of coast live oak in California. *Plant Disease* 97, 1025–1036.
- Makatini G.J., 2014. *The Role of Sucker Wounds as Portals for Grapevine Trunk Pathogen Infections*. MSc Thesis. University of Stellenbosch, South Africa, 108 pp.
- Marin-Felix Y., M. Hernandez-Restrepo, M.J. Wingfield, A. Akulov, A.J. Carnegie, R. Cheewangkoon, D. Gramaje, J.Z. Groenewald, V. Guarnaccia, F. Halleen, L. Lombaard, J. Luangsa-ard, S. Marincowitz, A. Moslemi, L. Mostert, W. Quaedvlieg, R.K. Schumacher, C.F.J. Spies, R. Thangavel, P.W.J. Taylor, A.M. Wilson, B.D. Wingfield, A.R. Wood and P.W. Crous, 2019. Genera of phytopathogenic fungi: GO-PHY 2. *Studies in Mycology* 92, 47–133.
- Mohammadi H. and Z. Banihashemi, 2012. First report of *Phaeoacremonium inflatipes* and *Phaeoacremonium mortoniae* associated with grapevine Petri disease in Iran. *Journal of Agricultural Science Techniques* 14, 1405–1414.
- Mohammadi H. and H. Hashemi, 2015. First report of *Phaeoacremonium aloesii* associated with grapevine Petri disease in Iran. *Journal of Plant Pathology* 97, 209–220.
- Mostert L., J.Z. Groenewald, R.C. Summerbell, V. Robert, D.A. Sutton, A.A. Padhye and P.W. Crous, 2005. Species of *Phaeoacremonium* associated with infections in humans and environmental reservoirs in infected woody plants. *Journal of Clinical Microbiology* 43, 1752–1767.
- Mostert L., F. Halleen, P.H. Fourie and P.W. Crous, 2006a. A review of *Phaeoacremonium* species involved in Petri disease and esca of grapevines. *Phytopathologia Mediterranea* 45, S12–S29.
- Mostert L., J.Z. Groenewald, R.C. Summerbell, W. Gams and P.W. Crous, 2006b. Taxonomy and pathology of *Togninia* (Diaporthales) and its *Phaeoacremonium* anamorphs. *Studies in Mycology* 54, 1–115.
- Moyo P., E. Allsopp, F. Roets, L. Mostert and F. Halleen, 2014. Arthropods vector grapevine trunk disease pathogens. *Phytopathology* 104, 1063–1069.
- Mugnai L., A. Graniti and G. Surico, 1999. Esca (black measles) and brown wood-streaking: two old and elusive diseases of grapevines. *Plant Disease* 83, 404–416.
- Munkvold G.P. and J.J. Marois, 1995. Factors associated with variation in susceptibility of grapevine pruning wounds to infection by *Eutypa lata*. *Phytopathology* 85, 249–256.
- Mutawila C., F. Halleen and L. Mostert, 2016. Optimisation of time of application of *Trichoderma* biocontrol agents for protection of grapevine pruning wounds. *Australian Journal of Grape and Wine Research* 22, 279–287.
- O'Donnell K. and E. Cigelnik, 1997. Two divergent intragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* 7, 103–116.
- Özben S., K. Değirmenci, F. Demirci and S. Uzunok, 2012. First report of *Phaeoacremonium scolyti* associated with esca and Petri disease of grapevine in Turkey. *Plant Disease* 96, 766.
- Quaglia M., L. Covarelli and A. Zizzerini, 2009. Epidemiological survey on esca disease in Umbria, central Italy. *Phytopathologia Mediterranea* 48, 84–91.
- Raimondo M.L., F. Lops and A. Carlucci, 2014. *Phaeoacremonium italicum* sp. nov., associated with esca of grapevine in southern Italy. *Mycologia* 106, 1119–1126.
- Rooney-Latham S.N., A. Eskalen and W.D. Gubler, 2005. Teleomorph formation of *Phaeoacremonium aleophilum* in Austria. *Plant Disease* 89, 177–184.
- Scheck H.J., S.J. Vasquez and W.D. Gubler, 1998. First report of the three *Phaeoacremonium* spp. causing young grapevine decline in California. *Plant Disease* 82, 590.
- Spies C.F.J., P. Moyo, F. Halleen and L. Mostert, 2018. *Phaeoacremonium* species diversity on woody hosts in the Western Cape Province of South Africa. *Persoonia* 40, 26–62.
- Úrbez-Torres J.R., P. Haag, P. Bowen and D.T. O'Gorman, 2014. Grapevine Trunk Diseases in British Columbia: Incidence and Characterization of the Fungal Pathogens Associated with Esca and Petri Diseases of Grapevine. *Plant Disease* 98, 469–482.
- Van Niekerk J.M., F. Halleen and P.H. Fourie, 2011. Temporal susceptibility of grapevine pruning wounds to trunk pathogen infection in South African grapevine. *Phytopathologia Mediterranea* 50, S139–S150.
- Weber E.A., F.P. Trouillas and W.D. Gubler, 2007. Double pruning of grapevines: A cultural practice to reduce infections by *Eutypa lata*. *American Journal of Enology and Viticulture* 58, 61–66.
- White C., F. Halleen and L. Mostert, 2011a. Symptoms and fungi associated with esca in South African vineyards. *Phytopathologia Mediterranea* 50, 236–246.
- White C., F. Halleen, M. Fischer and L. Mostert, 2011b. Characterization of the fungi associated with esca diseased grapevines in South Africa. *Phytopathologia Mediterranea* 50, 204–223.

Accepted for publication: December 26, 2018